Compression of Nakajima monomials in type A and C

Inaugural - Dissertation

zur

Erlangung des Doktorgrades der Mathematisch-Naturwissenschaftlichen Fakultät der Universität zu Köln

vorgelegt von

Matthias Meng

aus Sangerhausen

Gutachter Prof. Dr. P. Littelmann, Universität zu Köln (Betreuer)

Prof. Dr. H. Geiges, Universität zu Köln

Prof. Dr. A. Schilling, University of California, Riverside

Tag der Disputation 25.01.2011

Kurzzusammenfassung

Wir beschreiben für Lie-Algebren vom Typ A und C einen expliziten Kristallmorphismus zwischen Nakajima Monomen und Monomen, die eine Realisierung von kristallinen Basen von endlich dimensionalen irreduziblen Moduln der quantisierten universell einhüllenden Algebra bilden. Dieser Morphismus liefert eine Verbindung zwischen beliebigen Nakajima Monomen und Nakashima Kashiwara Tableaux, welche zu einer Übersetzung zwischen Nakajima Monomen und dem Wege-Modell von Littelmann führt. Außerdem definieren wir als eine Anwendung unserer Ergebnisse ein "Insertion scheme" für Nakajima Monome, das kompatibel mit dem "Insertion scheme" für Tableaux ist.

Abstract

We describe an explicit crystal morphism between Nakajima monomials and monomials which give a realization of crystal bases for finite dimensional irreducible modules over the quantized enveloping algebra for Lie algebras of type A and C. This morphism provides a connection between arbitrary Nakajima monomials and Nakashima Kashiwara tableaux. This yields a translation of Nakajima monomials to the Littelmann path model. Furthermore, as an application of our results we define an insertion scheme for Nakajima monomials compatible to the insertion scheme for tableaux.

Acknowledgements

I would like to thank Professor Peter Littelmann for inducting me into this subject and Professor Seok-Jin Kang for a short but inspiring discussion about my results.

This research was supported by the DFG-Graduiertenkolleg "Globale Strukturen in Geometrie und Analysis" Mathematisches Institut der Universität zu Köln, Weyertal 86-90, 50931 Köln, Germany.

Contents

Introduction		1	
1	Nakajima monomials		5
2	Typ	pe A	7
	2.1	Compression of Nakajima monomials in type A	7
	2.2	Monomials and A-Tableaux	20
	2.3	Insertion scheme for monomials in type A	21
3	Type C		26
	3.1	Compression of Nakajima monomials in type C	26
	3.2	Monomials and C -Tableaux	42
	3.3	Insertion scheme for monomials in type C	43
References		48	

Introduction

Crystal basis theory for integrable modules over quantum groups as introduced by Kashiwara [4] leads to a combinatorial interpretation of those modules in terms of crystals themselves, and furthermore their various models. Let us list some of those models which will play a role in the present thesis:

- 1. semistandard Young tableaux and reversed Young tableaux, satisfying certain conditions, for classical Lie algebras by Kashiwara and Nakashima [6], and Kim and Shin [7] (see also Kang, Kim, and Shin in [2] and [3]), respectively,
- 2. Young walls for affine Lie algebras by Kang, Kim, and Lee [1],
- 3. monomials for Kac-Moody algebras discovered by Nakajima [11], and generalized by Kashiwara [5],
- 4. the path model for symmetrizable Kac-Moody algebras introduced by Littelmann [10].

Let us be more precise about the monomial and the path model. Defining a t-analog of q-characters Nakajima [11] introduced a set of monomials \mathfrak{M} in certain variables $Y_i(k)$, and discovered a crystal structure on certain subsets of \mathfrak{M} . Kashiwara [5] generalized this, in that he defined a crystal structure on \mathfrak{M} , and proved that the connected component containing a highest weight monomial of integral weight λ is isomorphic to the crystal basis $B(\lambda)$ of irreducible highest weight modules.

Kang, Kim, and Shin [2], [3] considered specific highest weight monomials $M_{\lambda} \in \mathfrak{M}$ of weight λ and gave an explicit description of their connected components $\mathcal{M}(\lambda)$. Furthermore, they exhibited a connection between those and reversed Young tableaux.

As a generalization of Young tableaux Littelmann [9] considered paths (modulo reparametrization) on the real form of the weight lattice and defined the so-called root operators acting on those paths. With these operators the set of paths Π becomes a crystal, and every Young tableau can easily be considered as such a path [10].

In this thesis we describe a translation between the monomial and the path model. That is, we map an arbitrary monomial, not necessarily contained in some $\mathcal{M}(\lambda)$, to a path in Π such that our mapping yields a crystal morphism. For example, if the underlying Lie algebra is of type A_1 a possible definition of such a map is quite obvious: each monomial $M \in \mathfrak{M}$ is of the form M =

 $Y_1(i_1)^{y(i_1)} \cdots Y_1(i_k)^{y(i_k)}$ where $k \in \mathbb{N}$, $i_1, \ldots, i_k \in \mathbb{Z}$ and $i_1 < \ldots < i_k$, and $y(i_j) \in \mathbb{Z}$. To a fixed monomial M we associate the path $\pi_M = \pi_{y(i_k)\Lambda_1} * \ldots * \pi_{y(i_1)\Lambda_1}$, where $\pi_{\lambda}(t) = t\lambda$ is the path connecting the origin to λ .

Example. Consider the monomial $M = Y_1(2)^{-1}Y_1(1)^2$:


$$M = Y_1(2)^{-1}Y_1(1)^2 \qquad \longleftrightarrow \qquad \pi \qquad \begin{vmatrix} -\alpha_1 & -\Lambda_1 & 0 & \Lambda_1 & \alpha_1 \\ & & & & & \end{vmatrix}$$

$$\downarrow \tilde{f}_1 \qquad \qquad \downarrow \tilde{f}_1 \qquad \qquad \downarrow \tilde{f}_1$$

$$\tilde{f}_1 M = Y_1(2)^{-2}Y_1(1)^1 \qquad \longleftrightarrow \qquad \tilde{f}_1(\pi) \qquad \qquad \uparrow \tilde{f}_1(\pi) \qquad \qquad \downarrow \tilde{f}_1(\pi)$$

Note that, even for type A_2 , to find such a mapping is by far less obvious.

Example. For \mathfrak{g} of type A_2 we have $\tilde{f}_1(Y_1(2)^{-1}Y_1(1)^2) = Y_1(2)^{-2}Y_1(1)Y_2(1)$. Adopting the (obvious) construction in type A_1 , we would associate the path π displayed on the left:



After applying \tilde{f}_1 , observe that $\tilde{f}_1(\pi)$ has a linear part different from any fundamental root direction. That is, $\tilde{f}_1(\pi)$ does not coincide with the path we would associate to the monomial $\tilde{f}_1(M)$ in the same manner.

By generalizing the results of [2] and [3] in type A and C to arbitrary monomials in \mathfrak{M} , we determine the structure of the crystal graph associated to the connected component of an arbitrary, not necessarily highest weight, monomial in \mathfrak{M} . More precise, we give a crystal morphism between the set \mathfrak{M} and the set of tableaux which give realizations of $B(\lambda)$, and consequently, due to Littelmann [10], we can associate a path to those tableaux. Our crystal morphism compresses an arbitrary Nakajima monomial $M \in \mathfrak{M}$ into one which lies in a connected component $\mathcal{M}(\lambda)$, with integral dominant weight λ depending on M.

In a first step we describe a crystal isomorphism between the Nakajima monomials and certain matrices, namely $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$ in the A_n -case and $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$ in type C_n . This bijection allows us to define the compression of a monomial by compressing its associated matrix as follows: For simplicity let M denote the matrix associated to an arbitrary monomial $M \in \mathfrak{M}$ lying in some a priori unknown connected component of \mathfrak{M} . We give an algorithm which decomposes M into a sum $M = M_1 + M_2$, such that M_1 corresponds to a monomial in some $\mathcal{M}(\mu_1)$. Then, we move every column of M_2 one step to the left and denote by $M^{(1)}$ the sum of M_1 and the altered counterpart of M_2 . Our procedure allows an iteration yielding a sequence of matrices $M^{(i)}$. Since M has just finitely many nonzero columns, it is guaranteed that after a finite number of steps our iteration becomes stationary and we obtain a matrix $M^{(k)}$ corresponding to a monomial that lies in some $\mathcal{M}(\mu_k)$. We call $M^{(k)}$ the

compressed version of M. Our algorithm respects the crystal structure, that is we prove:

Main Theorem. Let \mathfrak{g} be of type A or C, and let $M \in \mathfrak{M}$ be a Nakajima monomial. Denote by $M^{(k)}$ its compressed version. Then, the map

$$\kappa: \mathfrak{M} \to \bigcup_{\lambda} \mathcal{M}(\lambda)$$

$$M \mapsto M^{(k)}$$


is a morphism of crystals. In particular, the connected component of M is isomorphic to the connected component of $\kappa(M)$.

Due to [2] and [3] we can assign a tableau S(N) to each $N \in \mathcal{M}(\lambda)$. Consequently, our Main Theorem gives:

Corollary. Let \mathfrak{g} be of type A or C, and let $M \in \mathfrak{M}$ be a Nakajima monomial. The mapping sending M to the tableau $S(\kappa(M))$ yields a crystal morphism.

Note that, in view of [10] we obtain a translation of Nakajima monomials into Littelmann paths.

Example. Consider the monomial $M = Y_1(2)Y_1(1)^2$, and the path obtained via our construction. Observe that our assignment commutes with the crystal operator \tilde{f}_1 , as illustrated in the following pictures.



As another application of our compression and the Corollary we define an insertion scheme for Nakajima monomials compatible with the insertion scheme of reversed tableaux described in [8]. More precise, let M_1 and M_2 be two matrices which correspond to arbitrary monomials in \mathfrak{M} . Then, we consider the matrix $M_1 * M_2 = (M_2, \mathbf{0}, M_1)$ with a suitable zero-matrix $\mathbf{0}$ and apply our compression procedure to $M_1 * M_2$. Following the convention that $M_1 * M_2$ interchangebly denotes the matrix and its associated monomial, we obtain $\kappa(M_1 * M_2) \in \bigcup_{\lambda} M(\lambda)$ and the tensor product rule of crystals yields

Theorem. Let \mathfrak{g} be of type A or C, and let \mathfrak{M} be the set of Nakajima monomials. Then, the map

$$\mathfrak{M} \otimes \mathfrak{M} \quad \to \quad \bigcup_{\lambda} \mathcal{M}(\lambda)$$
 $M_1 \otimes M_2 \quad \mapsto \quad \kappa(M_1 * M_2)$

is a morphism of crystals. In particular, the connected component of $M_1 \otimes M_2$ is isomorphic to the connected component of $\kappa(M_1 * M_2)$.

Chapter 1

Nakajima monomials

In this chapter we define the Nakajima monomials and their crystal structure. Let \mathfrak{g} be an arbitrary symmetrizable Kac-Moody Lie algebra with weight lattice P and I an index set such that $\alpha_i \in P$ for $i \in I$ are the simple roots. Let further $h_i \in P^*$ be the simple coroots and $(\cdot, \cdot) : P \times P \to \mathbb{Q}$ a bilinear symmetric form. For $i \in I$ and $\lambda \in P$ set $\langle h_i, \lambda \rangle := \frac{2(\alpha_i, \lambda)}{(\alpha_i, \alpha_i)}$.

For $i \in I$ and $n \in \mathbb{Z}$ we consider monomials in the variables $Y_i(n)$. That means we obtain the set of Nakajima monomials \mathfrak{M} as follows

$$\mathfrak{M}:=\bigg\{\prod_{i\in I,n\in\mathbb{Z}}Y_i(n)^{y_i(n)};y_i(n)\in\mathbb{Z} \text{ vanish except finitely many } (i,n)\bigg\}.$$

In order to define the crystal structure on \mathfrak{M} we take some integers $c = (c_{ij})_{i \neq j \in I} \subset \mathbb{Z}$ such that $c_{ij} + c_{ji} = 1$ and consider the monomials

$$A_i(n) := Y_i(n)Y_i(n+1) \prod_{j \neq i} Y_j(n+c_{ji})^{\langle h_j, \alpha_i \rangle}.$$

Let now M be an arbitrary monomial in \mathfrak{M} and $i \in I$. Then we set:

$$\begin{split} wt(M) &= \sum_{i} (\sum_{n} y_{i}(n)) \Lambda_{i}, \\ \varphi_{i}(M) &= \max\{ \sum_{k \leq n} y_{i}(k); n \in \mathbb{Z} \}, \\ \epsilon_{i}(M) &= \max\{ -\sum_{k > n} y_{i}(k); n \in \mathbb{Z} \}, \end{split}$$

where $\Lambda_i \in P$ are the fundamental weights, that means $\langle h_j, \Lambda_i \rangle = \delta_{i,j}$. To define the operators \tilde{e}_i and \tilde{f}_i we consider the values

$$n_f = \min\{n; \varphi_i(M) = \sum_{k \le n} y_i(k)\}$$
$$= \min\{n; \epsilon_i(M) = -\sum_{k > n} y_i(k)\},$$

$$n_e = \max\{n; \varphi_i(M) = \sum_{k \le n} y_i(k)\}$$
$$= \max\{n; \epsilon_i(M) = -\sum_{k > n} y_i(k)\}$$

and set

$$\tilde{f}_i(M) = \begin{cases} 0 & \text{if } \varphi_i(M) = 0, \\ A_i(n_f)^{-1}M & \text{if } \varphi_i(M) > 0, \end{cases}$$

$$\tilde{e}_i(M) = \begin{cases} 0 & \text{if } \epsilon_i(M) = 0, \\ A_i(n_e)M & \text{if } \epsilon_i(M) > 0. \end{cases}$$

Proposition 1.1 [5] With the maps $wt, \varphi_i, \epsilon_i, \tilde{f}_i$ and \tilde{e}_i thus defined, \mathfrak{M} becomes a semi-normal crystal.

We denote this crystal by \mathfrak{M}_c because the crystal structure of \mathfrak{M} depends on the choice of c. On the other hand one can easily see that the isomorphism class of the crystal \mathfrak{M}_c does not depend on this choice.

From now on, for simplicity, we choose $c = (c_{ij})_{i \neq j \in I}$ as follows:

$$c_{ij} = \begin{cases} 0 & \text{if } i > j, \\ 1 & \text{else.} \end{cases}$$

Now we recall the following result of Kashiwara.

Proposition 1.2 [5] Let M be a monomial of weight λ with $\tilde{e}_i(M) = 0$ for all $i \in I$. Then the connected component of \mathfrak{M} containing M is isomorphic to $B(\lambda)$.

The aim of this thesis is to give such an isomorphism explicitly for not necessarily highest weight monomials. In the first part we define this isomorphism for Lie algebras of type A. In the second part we generalize this to type C.

Chapter 2

Type A

2.1 Compression of Nakajima monomials in type A

Henceforth we consider a Lie algebra \mathfrak{g} of type A_n . In this case we have the fundamental weights $\Lambda_1, \ldots, \Lambda_n$ and we get an orthogonal basis $\beta_1, \ldots, \beta_{n+1}$ with $\beta_1 = \Lambda_1, \beta_i = \Lambda_i - \Lambda_{i-1}$ for $2 \le i \le n$ and $\beta_{n+1} = -\Lambda_n$. Moreover the simple roots are given by $\alpha_i = \beta_i - \beta_{i+1}$. Thus we compute

$$A_i(j) = Y_i(j)Y_i(j+1)Y_{i-1}(j+1)^{-1}Y_{i+1}(j)^{-1}.$$

For $i \in \{1, ..., n+1\}$ and $j \in \mathbb{Z}$ we introduce some specific monomials which will be of special interest to us:

$$X_i(j) := Y_{i-1}(j+1)^{-1}Y_i(j),$$

where we set $Y_{n+1}(j) = 1 = Y_0(j)$ for all $j \in \mathbb{Z}$.

With this notation we observe:

$$A_i(j) = X_i(j)X_{i+1}(j)^{-1}.$$

Let us shortly recall the monomial realization of the crystal bases $B(\lambda)$ given in [2]:

Proposition 2.1 [2] Let $\lambda = \sum_{k=1}^{n} a_k \Lambda_k$ be a dominant integral weight and consider $M_1 = Y_1(1)^{a_1} Y_2(1)^{a_2} \dots Y_n(1)^{a_n}$ as highest weight monomial. Then the connected component $\mathcal{M}_1(\lambda)$ of \mathfrak{M} containing M_1 is characterized as the set of monomials of the form

$$M = \prod_{\substack{i \in \{1, \dots, n+1\} \\ j \in \{1, \dots, n\}}} X_i(j)^{m_{ij}}$$

with

(i)
$$\sum_{i=1}^{n+1} m_{ij} = a_{j+1} + \ldots + a_n \text{ for } j = 1, \ldots, n,$$

(ii)
$$\sum_{k=i}^{n+1} m_{k,j} \le \sum_{k=i+1}^{n+1} m_{k,j-1}$$
 for $j = 2, \dots, n+1$ and $i = 1, \dots, n+1$.

For $s \in \mathbb{Z}$ we also consider the following *shifted* highest weight monomials of weight λ

$$M_s = Y_1(s)^{a_1} Y_2(s)^{a_2} \dots Y_n(s)^{a_n}.$$

As an immediate consequence of Proposition 2.1 we obtain their connected component $\mathcal{M}_s(\lambda)$ by the set of monomials of the form

$$M = \prod_{\substack{i \in \{1, \dots, n+1\}\\ j \in \{s, \dots, s+n-1\}}} X_i(j)^{m_{ij}}$$

satisfying condition (i) for $j = s, s+1, \ldots, s+n-1$ and (ii) for $i = 1, \ldots, n+1$ and $j = s+1, \ldots, s+n$.

Our aim is to *compress* an arbitrary monomial into the form of those in $\mathcal{M}_s(\lambda)$ for a suitable $\lambda \in P$ and $s \in \mathbb{Z}$ such that the crystal structure is preserved. As a first step we write an monomial in \mathfrak{M} as a product of $X_i(j)$'s. Thus we show that \mathfrak{M} is generated by the variables $X_i(j)$. That means we consider \mathfrak{M} as a group with the multiplication of monomials as binary operation. Let \mathbf{M} be the free abelian monoid generated by the set $\{X_i(j), i \in \{1, \ldots, n+1\}, j \in \mathbb{Z}\}$, with the same operation and we define an ideal $\mathbf{J} \subset \mathbf{M}$ by

$$\mathbf{J} = \langle \prod_{k=1}^{n+1} X_k (j+i-k), \text{ for } i = 1, \dots, n+1 \text{ and } j \in \mathbb{Z} \rangle_{\mathbf{M}}.$$

The quotient \mathbf{M}/\mathbf{J} becomes a group since we obtain the inverse of $X_i(j)$ by $\prod_{k=i+1}^{n+1} X_k(j-k+i) \prod_{k=1}^{i-1} X_k(j+i-k).$ Moreover we get

Proposition 2.2 Sending $X_i(j)$ onto $Y_i(j)Y_{i-1}(j+1)^{-1}$ yields a group isomorphism and therefore we get

$$\mathfrak{M} \cong \mathbf{M}/\mathbf{J}$$
.

Proof: In order to show surjectivity let M be of the form $M = \prod_{i \in I, j \in \mathbb{Z}} Y_i(j)^{y_i(j)}$.

First we write every $Y_{i-1}(j+1)^{-1}Y_i(j)$ that already occurs in M as $X_i(j)$. Then we consider the other $Y_i(j)^{y_i(j)}$'s in M. There are two possible cases: 1. case: $y_i(j) > 0$. Then we write

$$Y_{i}(j) = \prod_{k=0}^{i-1} Y_{k}(j+i-k)^{-1} \prod_{k=1}^{i} Y_{k}(j+i-k)$$
$$= \prod_{k=1}^{i} X_{k}(j+i-k).$$

Therefore we get

$$Y_i(j)^{y_i(j)} = \prod_{k=1}^i X_k(j+i-k)^{y_i(j)}.$$

2. case: $y_i(j) < 0$. In this case we get

$$Y_{i}(j)^{-1} = \prod_{\substack{k=i\\n+1}}^{n} Y_{k}(j-(k-i))^{-1} \prod_{\substack{k=i+1}}^{n+1} Y_{k}(j-(k-i))$$
$$= \prod_{\substack{k=i+1}}^{n+1} X_{k}(j-k+i).$$

Hence we have

$$Y_i(j)^{y_i(j)} = \prod_{k=i+1}^{n+1} X_k(j-k+i)^{-y_i(j)}.$$

These equations imply

$$Y_i(j)^{-1}Y_{i-1}(j+1) = \prod_{k=i+1}^{n+1} X_k(j-k+i) \prod_{k=1}^{i-1} X_k(j+i-k)$$

and hence with the definition of J we have injectivity.

Let now $M \in \mathfrak{M}$ be an arbitrary monomial. Due to Proposition 2.1 we can write M as a product of $X_i(j)$'s. That means we find $m_{ij} \in \mathbb{Z}_{>0}$ such that

П

$$M = \prod_{i \in \{1, \dots, n+1\}, j \in \mathbb{Z}} X_i(j)^{m_{ij}}.$$

Writing M in this way is obviously not unique. But we can fix a reduced notation $[m_{i,j}]$ and associate this matrix. Let us define the reduced notation on the level of matrices.

Let $M = m_{ij}$ be an arbitrary matrix in $\operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$, where $\operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ is the set of matrices with infinitely many columns but just finitely many different from zero. Then we get the reduced form [M] of M by applying the following rule:

(A1) For every $i \in \{1, ..., n+1\}$ we search for $j \in \mathbb{Z}$ such that

$$m_{i+s,i-s} \neq 0$$
 for all $s = -i+1, -i+2, \ldots, -1, 0, 1, \ldots, n-i$,

then we decrease these entries by

$$\min\{m_{i+s,j-s}; s=-i+1,-i+2,\ldots,-1,0,1,\ldots,n-i\}.$$

Denote by [M] the matrix obtained from M by applying this rule.

From now on we associate a matrix to a monomial in the following way: We write every $Y_i(j)^{y_i(j)}$ as a product of $X_k(l)$'s as in Proposition 2.1 and get a corresponding matrix $M = m_{ij}$. Then we apply (A1) and obtain [M]. We define an equivalence relation on $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$ by

$$M \sim N \text{ iff } [M] = [N]$$

and consider the quotient

$$\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim.$$

Now it is obvious that two matrices $M \sim N$ correspond to the same monomial and therefore we get a well defined map by sending a monomial to the associated matrix [M] as above. Moreover we obtain

Proposition 2.3 There exists a bijection between \mathfrak{M} and $Mat_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$.

Example 2.1 For \mathfrak{g} of type A_4 we consider the monomial

$$M = Y_1(4)^{-1}Y_3(1)Y_1(3)^{-1}Y_4(1)^{-1}Y_2(0)^2Y_3(2)^2.$$

Due to Proposition 2.1 we write:

$$Y_{1}(4)^{-1} = X_{2}(3)X_{3}(2)X_{4}(1)X_{5}(0),$$

$$Y_{3}(1) = X_{3}(1)X_{2}(2)X_{1}(1),$$

$$Y_{1}(3)^{-1} = X_{2}(2)X_{3}(1)X_{4}(0)X_{5}(-1),$$

$$Y_{4}(1)^{-1} = X_{5}(0),$$

$$Y_{2}(0)^{2} = X_{2}(0)^{2}X_{1}(1)^{2},$$

$$Y_{3}(2)^{2} = X_{3}(2)^{2}X_{2}(3)^{2}X_{1}(4)^{2}.$$

That means we obtain the associated reduced matrix by

$$\begin{pmatrix} 0 & 0 & 2 & 0 & 1 & 2 \\ 0 & 2 & 0 & 2 & 3 & 0 \\ 0 & 0 & 2 & 3 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 \end{pmatrix} \stackrel{(A1)}{=} \begin{pmatrix} 0 & 2 & 0 & 0 & 1 \\ 2 & 0 & 1 & 2 & 0 \\ 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix} = [M],$$

where we always only consider the finite part of the matrix which is different from zero.

Since we want this bijection to become a crystal morphism we need to define a crystal structure on $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ which coincides with the structure on \mathfrak{M} under our bijection.

Let
$$M = (m_{ij})_{i=1,\dots,n+1} \subset \mathbb{Z}_{\geq 0}$$
 be a $(n+1) \times \mathbb{Z}$ -matrix.
Set

$$wt(M) = \sum_{i=1}^{n+1} (\sum_{j \in \mathbb{Z}} m_{ij}) \beta_i,$$

$$\varphi_i(M) = \max \{ \sum_{j \le k} m_{ij} - \sum_{j < k} m_{i+1,j} ; k \in \mathbb{Z} \},$$

$$\epsilon_i(M) = -\min \{ \sum_{j > k} m_{ij} - \sum_{j \ge k} m_{i+1,j} ; k \in \mathbb{Z} \}.$$

If $\varphi_i(M) = 0$ we set $\tilde{f}_i(M) = 0$. Otherwise let $k \in \mathbb{Z}$ be minimal such that

$$\varphi_i(M) = \sum_{j \le k} m_{ij} - \sum_{j < k} m_{i+1,j}.$$

Note that, this k exists because M has just finitely many columns different from zero.

We define $\tilde{f}_i(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{i+1,k}$ (resp. $m_{i,k}$) by one. Formally spoken we get $\tilde{f}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(i,k), (i+1,k)\}, \\ m_{i,k} - 1 & \text{if } (s,j) = (i,k), \\ m_{i+1,k} + 1 & \text{if } (s,j) = (i+1,k). \end{cases}$$

Similarly, we can define the operator \tilde{e}_i :

If $\epsilon_i(M) = 0$ we set $\tilde{e}_i(M) = 0$.

For $\epsilon_i(M) \neq 0$ let $p \in \mathbb{Z}$ be maximal such that

$$\epsilon_i(M) = -(\sum_{j>p} m_{ij} - \sum_{j\geq p} m_{i+1,j}).$$

Then we obtain $\tilde{e}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(i,p), (i+1,p)\} ,\\ m_{i,p} + 1 & \text{if } (s,j) = (i,p),\\ m_{i+1,p} - 1 & \text{if } (s,j) = (i+1,p). \end{cases}$$

It is easy to see that these maps are well defined and that $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ along with $wt, \varphi_i, \epsilon_i, \tilde{f}_i$ and \tilde{e}_i becomes a semi-normal crystal. Now we can prove:

Proposition 2.4 The bijection

$$\Psi: \qquad \mathfrak{M} \qquad \to Mat_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$$

$$M = \prod_{i\in\{1,\dots,n+1\},j\in\mathbb{Z}} X_i(j)^{m_{ij}} \qquad \mapsto [m_{ij}]$$

is a crystal isomorphism.

Proof: We have to verify that for every $M \in \mathfrak{M}$ and $i \in I$ the following holds:

$$wt(M) = wt(\Psi(M)),$$

$$\varphi_i(M) = \varphi_i(\Psi(M)),$$

$$\epsilon_i(M) = \epsilon_i(\Psi(M)),$$

$$\Psi(\tilde{f}_i(M)) = \tilde{f}_i(\Psi(M)),$$

$$\Psi(\tilde{e}_i(M)) = \tilde{e}_i(\Psi(M)).$$

So let $M=\prod_{s\in I, t\in \mathbb{Z}}Y_s(t)^{y_s(t)}\in \mathfrak{M}$ be arbitrary and $\Psi(M)\in \mathrm{Mat}_{n+1\times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$

its corresponding reduced matrix. Now we show that

$$wt(M) = wt(\Psi(M)).$$

Assume we write M as a product of $X_s(t)$'s by writing every factor $Y_s(t)^{y_s(t)}$ as in Proposition 2.1 with corresponding matrix $m_{s,t}$. Now it suffices to show that wt(M) coincides with $wt(m_{s,t})$ because it is obvious that wt is invariant under (A1). So we get

$$wt(M) = \sum_{s} (\sum_{t} y_{s}(t)) \Lambda_{s}$$

$$= \sum_{s} (\sum_{t} y_{s}(t)) \beta_{1} + \ldots + \beta_{s}$$

$$= \sum_{s} (\sum_{t} (\sum_{j \leq s} y_{s}(t))) \beta_{s}$$

$$= \sum_{s} (\sum_{t} m_{st}) \beta_{s}$$

$$= \sum_{s} (\sum_{t} m_{st}) \beta_{s}$$
since wt is invariant under (A1),
$$= wt(\Psi(M)).$$

The same computations work for φ_i and ϵ_i .

Now we show that Ψ commutes with \tilde{f}_i : Let n_f be minimal such that $\varphi_i(M) = \sum_{t \leq n_f} y_s(t)$. Then we get $\tilde{f}_i(M) = A_i(n_f)^{-1}M$. Due to the choice of n_f we know that n_f is minimal such that

$$\varphi_i(M) = \varphi_i(\Psi(M)) = \sum_{j \le n_f} m_{ij} - \sum_{j < n_f} m_{i+1,j}.$$

That means we decrease (resp. increase) m_{i,n_f} (resp. m_{i+1,n_f}) by one in $\Psi(M)$.

But since $A_i(n_f)^{-1} = X_i(n_f)^{-1}X_{i+1}(n_f)$ it follows that $\tilde{f}_i(\Psi(M))$ is a corresponding matrix of $\tilde{f}_i(M)$. It remains to show that $\tilde{f}_i(\Psi(M)) = [\tilde{f}_i(\Psi(M))]$. Assume we had to apply (A1) only after having operated with \tilde{f}_i but not before. That means we get a full diagonal $m_{i+1+s,n_f-s} \neq 0$ for all $s = -i+1, -i+2, \ldots, -1, 0, 1, \ldots, n-i$ after having increased m_{i+1,n_f} by one. But due to the choice of n_f we have $m_{i,n_f+1} < m_{i+1,n_f}$ since otherwise $\sum_{j \leq n_f} m_{ij} - \sum_{j < n_f} m_{i+1,j}$ would not be maximal. Therefore increasing m_{i+1,n_f} doesn't cause any new (A1) application and

$$\Psi(\tilde{f}_i(M)) = \tilde{f}_i(\Psi(M)).$$

The same arguments hold for \tilde{e}_i which finishes our proof.

Now we define the set of matrices such that the corresponding monomials give a realization of the crystal bases $B(\lambda)$.

Definition 2.1 Define $\mathfrak{N} \subset Mat_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ as the set of matrices whose reduced forms have only zero-entries out of an $(n+1) \times n$ -submatrix $M = (m_{ij})_{\substack{i=1,\ldots,n+1\\j=0,\ldots,n-1}}$ with the following properties:

(i)
$$m_{ij} \in \mathbb{Z}_{\geq 0}$$
 for $i = 1, ..., n + 1$ and $j = 0, ..., n - 1$.

(ii)
$$\sum_{k=i}^{n+1} m_{k,j} \le \sum_{k=i+1}^{n+1} m_{k,j-1}$$
 for $i = 1, \dots, n+1$ and $j = 1, \dots, n-1$,
where we set $\sum_{k=i+1}^{n+1} m_{k,j-1} = 0$ for $i = n+1$.

Due to Proposition 2.1 the associated monomials of matrices in \mathfrak{N} can be considered as elements in $\mathcal{M}_s(\lambda)$ for a suitable $\lambda \in P$ and $s \in \mathbb{Z}$. Hence instead of a crystal morphism between \mathfrak{M} and $\bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$ which we originally intended to find, we just need a morphism from $\mathrm{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ to \mathfrak{N}/\sim . The idea is to compress the matrices. More precisely we move entries into the next column to the left such that the crystal structure is preserved. We do this by decomposing our matrix $M = M_1 + M_2$ with $M_1 \in \mathfrak{N}$ according to the following rule:

The lower decomposition rule:

Let $M = [M] = m_{ij} \in \operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ be a reduced version of an arbitrary matrix. Let $k \in \mathbb{Z}$ be minimal and $l \in \mathbb{Z}$ be maximal such that $m_{ij} = 0$ for all j < l, j > k and $i \in \{1, \ldots, n+1\}$. That means the finite part of M which is different from zero is an $(n+1) \times (l-k+1)$ -matrix over $\mathbb{Z}_{\geq 0}$. For simplicity we set p = l - k and renumber the columns by $0, \ldots, p$. We also assume that $p \geq n-1$, otherwise we fill the matrix with zero-entries on the right side. We search for $M_1 \in \mathfrak{N}$ such that

$$M = M_1 + M_2.$$

We explain how to compute $M_1 = m_{ij}^{(1)}$ out of $M = m_{i,j}$ recursively:

For i = 1, ..., n + 1 we set $m_{i,0}^{(1)} := m_{i,0}$.

Then, for each j from 1 to p we do the following:

For i = n + 1 to i = 1 we compare

$$\sum_{k>i+1} m_{k,j-1}^{(1)} \text{ with } m_{i,j} + \sum_{k>n+1} m_{k,j}^{(1)}$$

and if $\sum_{k \ge i+1} m_{k,j-1}^{(1)} < m_{i,j} + \sum_{k \ge p+1} m_{k,j}^{(1)}$ then we set

$$m_{i,j}^{(1)} := \sum_{k \geq i+1} m_{k,j-1}^{(1)} - \sum_{k \geq i+1} m_{k,j}^{(1)}.$$

Otherwise, namely if $\sum_{k \geq i+1} m_{k,j-1}^{(1)} \geq m_{i,j} + \sum_{k \geq p+1} m_{k,j}^{(1)}$, we set

$$m_{i,j}^{(1)} = m_{i,j}.$$

This way we get M_1 and set

$$M_2 := M - M_1$$
.

By construction it is obvious that M_1 satisfies condition (i) and (ii) of Definition 2.1.1 but it remains to show that M_1 has at most n columns different from zero such that we can guarantee that $M_1 \in \mathfrak{N}$. For that we show

Lemma 2.1 Let $[M] = (m_{ij})_{\substack{i=1,\dots,n+1\\j=0,\dots l-1}} \in Mat_{n+1\times l}(\mathbb{Z}_{\geq 0})$ be a reduced matrix without zero columns which satisfies condition (ii) of Definition 2.1.1. Then we have

$$l < n$$
.

Proof: Since $\sum_{k=i+1}^{n+1} m_{k,0} = 0$ for i = n+1 condition (ii) yields $m_{n+1,1} = 0$. Again by condition (ii) we obtain $0 = m_{n+1,1} \ge m_{n,2} + m_{n+1,2}$ and therefore $m_{n,2} = m_{n+1,2} = 0$. In general condition (ii) provides

$$\sum_{k=n+2-j}^{n+1} m_{k,j} = 0.$$

That means in particular that $m_{i,n} = 0$ for all i = 2, ..., n + 1 and $m_{i,j} = 0$ for all $i \in I$ and $j \ge n + 1$.

It remains to show that $m_{1,n} = 0$.

Assume $m_{1,n} \neq 0$. Since $\sum_{k=n+2-j}^{n+1} m_{k,j} = 0$ this implies $m_{1+t,n-t} \geq m_{1,n}$ for all $t = 1, \ldots, n$. That means we can apply (A1) which is a contradiction to M being in reduced form and hence

$$m_{1,n} = 0.$$

Example 2.2 For \mathfrak{g} of type A_4 we consider the matrix

$$M = (m_{i,j})_{\substack{i=1,\dots,5\\j=0,\dots,4}} = \begin{pmatrix} 0 & 2 & 0 & 0 & 1\\ 2 & 0 & 1 & 2 & 0\\ 0 & 1 & 2 & 0 & 0\\ 0 & 0 & 0 & 0 & 0\\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Until we reach the entry $m_{3,2}$ we always get $\sum_{k\geq i+1} m_{k,j-1}^{(1)} \geq m_{i,j} + \sum_{k\geq p+1} m_{k,j}^{(1)}$ and set $m_{i,j}^{(1)} := m_{i,j}$, but at $m_{3,2}$ we have $\sum_{k\geq 4} m_{k,1}^{(1)} = 0 < 2 = \sum_{k\geq 4} m_{k,2}^{(1)} + m_{3,2}$. Therefore we set $m_{3,2}^{(1)} := \sum_{k\geq 4} m_{k,1}^{(1)} - \sum_{k\geq 4} m_{k,2}^{(1)} = 0$. The same happens once more at $m_{2,3}$ and hence

$$M_1 = egin{pmatrix} 0 & 2 & 0 & 0 & 1 \ 2 & 0 & 1 & 0 & 0 \ 0 & 1 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ 1 & 0 & 0 & 0 & 0 \end{pmatrix} \in \mathfrak{N}.$$

Moreover we obtain

Now we can define our desired map:

$$\Phi : \operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0}) / \sim \to \mathfrak{N} / \sim m_{ij} \mapsto n_{ij},$$

where the matrix n_{ij} is computed in the following way:

Let M be the reduced version of an arbitrary matrix in $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$. Then we consider the lower decomposition of M:

$$M = M_1 + M_2$$

with $M_1 \in \mathfrak{N}$.

Then we move every entry of M_2 one column to the left and denote the new M_2 by $M_2^{(1)}$. Now we set $M^{(1)} := M_1 + M_2^{(1)}$ and if $M^{(1)} \in \mathfrak{N}$ we are done and set

$$M^{(1)} =: N = n_{ij}$$
.

If $M^{(1)} \notin \mathfrak{N}$ we consider the lower decomposition of $[M^{(1)}]$ and do the same again. This iteration yields a sequence of matrices $M^{(i)}$. Since M has just finitely many columns different from zero there exists a $k \in \mathbb{N}$ such that the iteration becomes stationary with $M^{(k)} \in \mathfrak{N}$ and we set:

$$M^{(k)} =: N = n_{ij}.$$

Let us now combine Φ and Ψ to obtain the *compression map* κ from the set of arbitrary monomials into the set of monomials which give a realization of the crystal bases $B(\lambda)$:

$$\kappa := \Psi^{-1} \circ \Phi \circ \Psi : \mathfrak{M} \to \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda).$$

Before we prove that this map is a crystal morphism we continue our example:

Example 2.3 As above, let \mathfrak{g} be of type A_4 and take the monomial

$$M = Y_1(4)^{-1}Y_3(1)Y_1(3)^{-1}Y_4(1)^{-1}Y_2(0)^2Y_3(2)^2.$$

We have already seen that M corresponds to the reduced version matrix

$$\begin{pmatrix} 0 & 2 & 0 & 0 & 1 \\ 2 & 0 & 1 & 2 & 0 \\ 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

We obtained the lower decomposition of M by:

and therefore

$$M^{(1)} = \begin{pmatrix} 0 & 2 & 0 & 1 \\ 2 & 0 & 3 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

Since $M^{(1)} \notin \mathfrak{N}$ we decompose $[M^{(1)}] = M^{(1)}$ and get

$$M^{(2)} = egin{pmatrix} 0 & 2 & 0 & 1 \ 2 & 2 & 1 & 0 \ 2 & 1 & 0 & 0 \ 0 & 0 & 0 & 0 \ 1 & 0 & 0 & 0 \end{pmatrix} \in \mathfrak{N}$$

Now we apply Ψ^{-1} and receive the monomial

$$N = \Psi^{(-1)}(M^{(2)}) = X_1(1)^2 X_1(3) X_2(0)^2 X_2(1) X_3(0)^2 X_3(1) X_5(0)$$

= $Y_1(3) Y_2(0)^2 Y_1(2)^{-2} Y_3(0)^2 Y_3(1) Y_4(1)^{-1}$.

Due to Proposition 2.1 we get: $N \in \mathcal{M}_0(4\Lambda_2 + \Lambda_4)$.

Theorem 2.1 Let g be of type A. Then, the map

$$\kappa : \mathfrak{M} \to \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$$

$$M \mapsto (\Psi^{-1} \circ \Phi \circ \Psi)(M)$$

defined as above is a morphism of crystals.

Proof: We have already seen that Ψ is a crystal morphism therefore we limit the proof to Φ and get the claim by composition. Since Φ is successively defined it suffices to show that sending a reduced version matrix M to $M^{(1)}$ preserves the crystal structure. So we take such a matrix $M = m_{ij}$ and its lower decomposition $M = M_1 + M_2$ with $M_1 = m_{ij}^{(1)}$ and $M_2 = m_{ij}^{(2)}$.

By definition we obtain

$$(M^{(1)})_{ij} = m_{ij}^{(1)} + m_{i,j+1}^{(2)}.$$

Now we have to show the following for $i \in I$:

- (i) $wt(M) = wt(M^{(1)}),$
- (ii) $\varphi_i(M) = \varphi_i(M^{(1)}),$
- (iii) $\epsilon_i(M) = \epsilon_i(M^{(1)})$

and that computing $M^{(1)}$ interchanges with the Kashiwara operators, namely

(iv)
$$(\tilde{f}_i(M))^{(1)} = \tilde{f}_i(M^{(1)}),$$

(v)
$$(\tilde{e}_i(M))^{(1)} = \tilde{e}_i(M^{(1)}).$$

Since $\sum_{j\in\mathbb{Z}} m_{ij}$ doesn't change it is obvious that wt is invariant under this construction.

We prove (ii) and (iv) simultaneously and (iii) and (v) follow in an analogous manner. For simplicity we set $M^{(1)} =: N = n_{ij}$. We know that

$$\varphi_i(M) = \max\{\sum_{i \le k} m_{ij} - \sum_{i \le k} m_{i+1,j}; k \in \mathbb{Z}\}.$$

Assume $\varphi_i(M) \neq 0$ and let $k \in \mathbb{Z}$ be minimal such that

$$\varphi_i(M) = \sum_{j \le k} m_{ij} - \sum_{j < k} m_{i+1,j}.$$

Now assume $\varphi_i(M) < \varphi_i(N)$.

That is only possible if there exists $p \in \mathbb{Z}$ such that $m_{i,p+1}^{(2)} > m_{i+1,p}^{(2)}$. Otherwise $\sum_{j \leq p} n_{ij} - \sum_{j < p} n_{i+1,j}$ is equal to or smaller than $\sum_{j \leq p} m_{ij} - \sum_{j < p} m_{i+1,j}$. Due to lower decomposition this implies

$$(2.1) m_{i,p+1}^{(2)} \le m_{i,p+1} - m_{i+1,p}^{(1)}$$

and since $M = M_1 + M_2$,

$$(2.2) m_{i+1,p}^{(2)} = m_{i+1,p} - m_{i+1,p}^{(1)}$$

Now we compute

$$\begin{split} \sum_{j \leq p} n_{ij} - \sum_{j < p} n_{i+1,j} &= \sum_{j \leq p} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} - (\sum_{j < p} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)}) \\ &= \sum_{j < p} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,p}^{(1)} + m_{i,p+1}^{(2)} \\ &- (\sum_{j < p-1} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)} + m_{i+1,p-1}^{(1)} + m_{i+1,p}^{(2)}) \\ &\leq \sum_{j < p} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,p}^{(1)} + m_{i,p+1} - m_{i+1,p}^{(1)} \\ &- (\sum_{j < p-1} m_{i+1,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i+1,j+1}^{(1)} + m_{i+1,p-1}^{(2)} + m_{i+1,p}^{(2)}) \\ &\stackrel{(2.2)}{=} \sum_{j < p} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,p}^{(1)} + m_{i,p+1} - m_{i+1,p}^{(1)} \\ &- (\sum_{j < p-1} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)} + m_{i+1,p+1}^{(1)} + m_{i+1,p-1}^{(1)} + m_{i+1,p-1}^{(1)} + m_{i+1,p}^{(1)}) \\ &= \sum_{j \leq p+1} m_{ij} - \sum_{j < p+1} m_{i+1,j}. \end{split}$$

But due to the choice of k we have

$$\varphi_i(M) \ge \sum_{j \le p+1} m_{ij} - \sum_{j < p+1} m_{i+1,j}$$

and hence

$$\varphi_i(M) \geq \varphi_i(N)$$
.

Now suppose $\varphi_i(N) < \varphi_i(M)$. It is obvious that $\sum_{j \le k} n_{ij} - \sum_{j < k} n_{i+1,j}$ becomes smaller than $\sum_{j \le k} m_{ij} - \sum_{j < k} m_{i+1,j}$ if and only if:

$$0 \neq m_{i+1,k}^{(2)} > m_{i,k}^{(2)}$$
.

Moreover the choice of k again implies that

$$m_{i+1,k}^{(1)} \le m_{i+1,k} < m_{i,k}.$$

That means we have the following situation in M_1 :

$$\sum_{l>i+1} m_{l,k}^{(1)} = \sum_{l>i+1} m_{l,k-1}^{(1)}$$

and therefore

$$m_{i,k}^{(2)} = m_{i,k} - m_{i+1,k-1}^{(1)}$$

With this equation we compute:

$$\begin{split} \sum_{j \leq k-1} n_{ij} - \sum_{j < k-1} n_{i+1,j} &= \sum_{j \leq k-1} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} - (\sum_{j < k-1} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)}) \\ &= \sum_{j < k-2} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,k-1}^{(1)} + m_{i,k}^{(2)} \\ &- (\sum_{j < k-2} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)} + m_{i+1,k-1}^{(1)} + m_{i+1,k-1}^{(1)} \\ &= \sum_{j < k-2} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,k-1}^{(1)} + m_{i,k} - m_{i+1,k-1}^{(1)} \\ &- (\sum_{j < k-2} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)} + m_{i+1,k-2}^{(1)} \\ &+ m_{i+1,k-1} - m_{i+1,k-1}^{(1)}) \\ &= \sum_{j < k-2} m_{i,j}^{(1)} + m_{i,j+1}^{(2)} + m_{i,j+1}^{(1)} + m_{i,k-1}^{(1)} + m_{i,k} \\ &- (\sum_{j < k-2} m_{i+1,j}^{(1)} + m_{i+1,j+1}^{(2)} + m_{i+1,k-2}^{(1)} \\ &+ m_{i+1,k-1}) \\ &= \sum_{j \leq k} m_{ij} - \sum_{j < k} m_{i+1,j}. \end{split}$$

Hence

$$\varphi_i(N) \ge \varphi_i(M)$$
.

Furthermore these computations also show that we obtain $\varphi_i(N)$ either by

$$\sum_{j \le k} n_{ij} - \sum_{j \le k} n_{i+1,j}$$

or as in the last case by

$$\sum_{j \le k-1} n_{ij} - \sum_{j < k-1} n_{i+1,j}$$

and that k (resp. k-1) is minimal with this property.

That means that we obtain $\tilde{f}_i(N)$ by operating on $n_{i,k}$ or on $n_{i,k-1}$. So assume we operate on $n_{i,k}$ and consider the lower decomposition of $\tilde{f}_i(M)$. In this case we know that $m_{i,k} > m_{i+1,k-1}$ and the same in N. That means $n_{i,k} > n_{i+1,k-1}$.

Hence

$$m_{i,k}^{(1)} > m_{i+1,k-1}^{(1)}$$

That implies

$$\sum_{j \geq i+2} m_{j,k-1}^{(1)} > \sum_{j \geq i+1} m_{j,k}^{(1)}$$

and in particular

$$m_{i+1,k}^{(2)} = 0.$$

Therefore we get $m_{i+1,k}^{(1)}$ increased by one in the lower decomposition of $\tilde{f}_i(M)$. Moreover we know that

$$m_{i+1,k}^{(1)} = m_{i+1,k} \ge m_{i,k+1}.$$

That means

$$m_{i,k+1}^{(2)} = 0$$

and increasing $m_{i+1,k}^{(1)}$ by one doesn't change the decomposition of the k+1-st column.

Hence

$$(\tilde{f}_i(M))^{(1)} = \tilde{f}_i(N).$$

Same arguments show that $m_{i+1,k}^{(2)}$ is increased by one if we operate on $n_{i,k-1}$ and we also get

$$(\tilde{f}_i(M))^{(1)} = \tilde{f}_i(N).$$

For $M \in \mathfrak{M}$ we set B(M) to be the connected component of M in \mathfrak{M} .

Corollary 2.1 For $M \in \mathfrak{M}$ and $\Phi(\Psi(M)) =: n_{i,j} \in \mathfrak{N}$ we consider $s \in \mathbb{Z}$ maximal such that $n_{i,j} = 0$ for all j < s and $i \in \{1, ..., n+1\}$. Furthermore for k = 1, ..., n+1 we define the values:

$$a_k := \sum_{i \in \{1, \dots, n+1\}} n_{i,k+s-1} - \sum_{i \in \{1, \dots, n+1\}} n_{i,k+s} \ge 0.$$

Then we have

$$\kappa(M) \in \mathcal{M}_s(\sum_{k=1}^n a_k \Lambda_k)$$

and hence by restriction

$$\kappa_{|B(M)}$$
 : $B(M) \rightarrow \mathcal{M}_s(\sum_{k=1}^n a_k \Lambda_k)$

is a crystal isomorphism.

2.2 Monomials and A-Tableaux

In this section we give an application of the compression defined in Section 2.1. In their framework about the correspondence between Young walls and Young tableaux, Kim and Shin [7] gave another realization of the crystal bases $B(\lambda)$ in the sense of reversed Young tableaux. Moreover Kang, Kim and Shin [2] constructed a crystal morphism between the monomials in $\mathcal{M}_1(\lambda)$ for dominant integral weights λ and those reversed tableaux. By combining this with the crystal morphism κ defined in Section 2.1 we can generalize their morphism to arbitrary monomials in \mathfrak{M} .

- **Definition 2.2** (i) We define a reversed Young diagram to be a collection of boxes in rightjustified rows with a weakly decreasing number of boxes in each row from bottom to top.
 - (ii) We define a (reversed) tableau by a reversed Young diagram filled with positive integers.
- (iii) A (reversed) tableau S is called a (reversed) semistandard tableau if the entries in S are weakly increasing from left to right in each row and strictly increasing from top to bottom in each column.

For a dominant integral weight λ we define $S(\lambda)$ to be the set of all (reversed) semistandard tableaux of shape λ with entries $1, \ldots, n+1$, which gives a realization of the crystal bases $B(\lambda)$ [7].

Let $M \in \mathcal{M}_1(\lambda)$ be a monomial and m_{ij} the associated reduced matrix in \mathfrak{N} . We define the tableau S(M) to be the semistandard reversed tableau with m_{ij} -many i entries in j-th row. Then we get

Proposition 2.5 [2] The map

$$\Omega : \mathcal{M}_1(\lambda) \to S(\lambda)
M \mapsto S(M)$$

is a crystal isomorphism.

It is obvious how to generalize this morphism to $\mathcal{M}_s(\lambda)$. Let $M \in \mathcal{M}_s(\lambda)$ be a monomial and m_{ij} the associated reduced matrix in \mathfrak{N} . In this case we define S(M) to be the semistandard reversed tableaux with m_{ij} -many i entries in the j-s+1-st row and get the morphism

$$\Omega : \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda) \to \bigcup_{\lambda \in P} S(\lambda)$$

$$M \mapsto S(M).$$

The combination of this morphism with the compression map κ yields:

Corollary 2.2 The map

$$\Omega\circ\kappa \ : \ \mathfrak{M} \ \to \ \bigcup_{\lambda\in P}S(\lambda)$$

is a crystal morphism.

Example 2.4 For \mathfrak{g} of type A_4 we consider and the monomial

$$M = Y_1(4)^{-1}Y_3(1)Y_1(3)^{-1}Y_4(1)^{-1}Y_2(0)^2Y_3(2)^2.$$

We have already seen that

$$\kappa(M) = N = X_1(1)^2 X_1(3) X_2(0)^2 X_2(1) X_3(0)^2 X_3(1) X_5(0),$$

with the corresponding reduced matrix

$$\begin{pmatrix} 0 & 2 & 0 & 1 \\ 2 & 2 & 1 & 0 \\ 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Therefore we assign the following semistandard reversed Young tableau

$$S(M) = \frac{\boxed{\frac{1}{2}}}{\boxed{\frac{1}{2} \mid 2 \mid 2 \mid 3 \mid 3 \mid 5}}.$$

2.3 Insertion scheme for monomials in type A

In this section, as another application of the compression given in Section 2.1, we define a bumping rule for Nakajima monomials. That means we consider the crystal tensor product of two monomials M_1 and M_2 and search for a monomial $N \in \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$ such that the connected component of $M_1 \otimes M_2$ is isomorphic to the connected component of N. Moreover we will see that this bumping is compatible with the reversed bumping for reversed tableaux (given in [8]).

Before we define the monomial bumping we recall the tensor product rule for crystals B_1 and B_2 :

The set $B_1 \otimes B_2 := \{b_1 \otimes b_2; b_1 \in B_1 \text{ and } b_2 \in B_2\}$ becomes a crystal by setting

$$wt(b_1 \otimes b_2) = wt(b_1) + wt(b_2),$$

$$\epsilon_i(b_1 \otimes b_2) = \max\{\epsilon_i(b_1), \epsilon_i(b_2) + \langle h_i, wt(b_1) \rangle\},$$

$$\varphi_i(b_1 \otimes b_2) = \max\{\varphi_i(b_1) + \langle h_i, wt(b_2) \rangle, \varphi_i(b_2)\},$$

$$\tilde{f}_i(b_1 \otimes b_2) = \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \epsilon_i(b_2), \\ b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \leq \epsilon_i(b_2), \end{cases}$$

$$\tilde{e}_i(b_1 \otimes b_2) = \begin{cases} b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \epsilon_i(b_2), \\ \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \leq \epsilon_i(b_2), \end{cases}$$

Let now M_1 and M_2 be reduced matrices of monomials in \mathfrak{M} . In order to use the compression procedure we associate a matrix $M_1 * M_2 \in \operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$

to the tensor product $M_1 \otimes M_2$ in the following way:

$$M_1 \otimes M_2 \mapsto \left(\begin{array}{ccc} & 0 & & & \\ & 0 & & & \\ & M_2 & \vdots & M_1 & \\ & 0 & & \\ & 0 & & \end{array} \right) =: M_1 * M_2,$$

where again M_1 and M_2 stand for their finite parts different from zero. With the crystal structure on $\operatorname{Mat}_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ we can show:

Proposition 2.6 The map

$$Mat_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \otimes Mat_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \rightarrow Mat_{n+1\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim M_1\otimes M_2 \mapsto M_1*M_2$$

is a crystal morphism.

Proof: Let M_1 and M_2 be matrices as above and after possible renumbering we set $M_1 = (m_{i,j}^1)_{\substack{i=1,\ldots,n+1\\j=1,\ldots,l}}$ and $M_2 = (m_{i,j}^2)_{\substack{i=1,\ldots,n+1\\j=1,\ldots,t}}$. For simplicity we write $M_1 * M_2 =: M = m_{i,j}$.

Since M_1 and M_2 are reduced we get by definition that $M_1 * M_2$ is a reduced matrix without any (A1) application.

We have to show that wt, φ_i and ϵ_i are invariant under this map and that it commutes with \tilde{f}_i and \tilde{e}_i . We observe directly from the definition that

$$wt(M_1 \otimes M_2) = wt(M_1 * M_2).$$

For $i \in I$ we show that

$$\varphi_i(M_1 \otimes M_2) = \varphi_i(M_1 * M_2).$$

In order to do this we distinguish the two cases $\varphi_i(M_1) > \epsilon_i(M_2)$ and $\varphi_i(M_1) \le \epsilon_i(M)$. At first we assume $\varphi_i(M_1) > \epsilon_i(M_2)$ and take k minimal such that

$$\varphi_i(M_1) = \sum_{j \le k} m_{i,j}^1 - m_{i+1,j-1}^1.$$

This implies

$$\varphi_i(M_1*M_2) = \sum_{j \leq k+t+1} m_{i,j} - m_{i+1,j-1}.$$

$$= \sum_{j \leq k} m_{i,j}^1 - m_{i+1,j-1}^1$$

$$+ \sum_j m_{i,j}^2 - \sum_j m_{i+1,j}^2$$
since $\langle h_i, \Lambda_j \rangle = \delta_{i,j}$,
$$= \varphi_i(M_1) + \sum_j m_{i,j}^2 \langle h_i, \Lambda_i - \Lambda_{i-1} \rangle$$

$$+ \sum_j m_{i+1,j}^2 \langle h_i, \Lambda_{i+1} - \Lambda_i \rangle$$

$$= \varphi_i(M_1) + \sum_j m_{i,j}^2 \langle h_i, \beta_i \rangle$$

$$+ \sum_j m_{i+1,j}^2 \langle h_i, \beta_{i+1} \rangle$$

$$= \varphi_i(M_1)$$

$$+ \left\langle h_i, \sum_j m_{i,j}^2 \beta_i + \sum_j m_{i+1,j}^2 \beta_{i+1} \right\rangle$$

$$\langle h_i, \beta_j \rangle = 0 \text{ for } j \neq i, i+1, \qquad = \varphi_i(M_1) + \left\langle h_i, \sum_i (\sum_j m_{i,j}^2) \beta_i \right\rangle$$

$$= \varphi_i(M_1) + \langle h_i, wt(M_2) \rangle$$
since $\varphi_i(M_1) > \epsilon_i(M_2)$,
$$= \varphi_i(M_1 \otimes M_2).$$

Moreover these computations also show: If $\varphi_i(M_1) > \epsilon_i(M_2)$, then k is minimal with

$$\varphi_i(M_1 * M_2) = \varphi_i(M) = \sum_{j \le k} m_{i,j} - \sum_{j < k} m_{i+1,j}$$

and hence

$$\tilde{f}_i(M_1 * M_2) = \tilde{f}_i M_1 * M_2.$$

Due to the tensor product rule we also observe

$$\tilde{f}_i(M_1 \otimes M_2) = \tilde{f}_i M_1 \otimes M_2,$$

which implies that the map interchanges with \tilde{f}_i in this case.

Let now $\varphi_i(M_1) \leq \epsilon_i(M_2)$. This yields directly $\varphi_i(M_1 * M_2) = \varphi_i(M_2) = \varphi_i(M_1 \otimes M_2)$ and again $\tilde{f}_i(M_1 * M_2) = M_1 * \tilde{f}_i M_2$. Same arguments hold for ϵ_i and \tilde{e}_i .

With this interpretation of the crystal tensor product of monomials we are able to give the definition of bumping for Nakjima monomials. Let M_1 and M_2 be monomials in \mathfrak{M} then we define $M_1 \to M_2$ as the result of the following compositions of crystal morphisms:

$$\mathfrak{M} \times \mathfrak{M} \longrightarrow \operatorname{Mat}_{n+1 \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \stackrel{\Phi}{\to} \mathfrak{N}/\sim \longrightarrow \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$$
$$(M_1, M_2) \mapsto \Psi(M_1) * \Psi(M_2) \stackrel{\Phi}{\mapsto} N \mapsto \Psi^{-1}(N).$$

In other words we set

$$M_1 \to M_2 := \Psi^{-1}(\Phi(\Psi(M_1) * \Psi(M_2))).$$

With Proposition 2.6 and Theorem 2.1 we observe

Theorem 2.2 Let g be of type A. Then, the map

$$\mathfrak{M} \otimes \mathfrak{M} \quad \to \quad \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$$
 $M_1 \otimes M_2 \quad \mapsto \quad M_1 \to M_2$

defined as above is a morphism of crystals.

Now we notice that the monomial bumping coincides with the tableaux bumping defined in [8]. More precisely, if we take $M_1, M_2 \in \mathfrak{M}$ we have two possibilities to associate a reversed tableaux to their tensor product $M_1 \otimes M_2$. The first one is to take the monomial bumping $M_1 \to M_2$ and to consider the tableaux $S(M_1 \to M_2)$ in the sense of Corollary 2.2. On the other hand we compute $S(M_1)$ and $S(M_2)$ and apply the reversed bumping rule given by Kim and Shin [7], [8], namely $S(M_1) \to S(M_2)$. Corollary 2.2 and Theorem 2.2 imply

$$S(M_1 \to M_2) = S(M_1) \to S(M_2).$$

In order to put this result in concrete terms we consider the following

Example 2.5 Let \mathfrak{g} be of type A_3 , say $\mathfrak{g} = \mathfrak{sl}_4$, and take

$$M_1 = Y_1(2)^{-1}Y_3(1)Y_2(1)Y_2(2)^{-1},$$

 $M_2 = Y_2(3)^{-2}Y_3(2)^{-1}Y_1(1)^3Y_1(3).$

As in section 2 we assign the matrix to M_1 by

$$Y_1(2)^{-1} = X_2(1)X_3(0)X_4(-1),$$

$$Y_3(1) = X_3(1)X_2(2)X_1(3),$$

$$Y_2(1) = X_2(1)X_1(2),$$

$$Y_2(2)^{-1} = X_3(1)X_4(0).$$

Therefore

$$M_1 \leftrightarrow \begin{pmatrix} 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{pmatrix} \stackrel{(A1)}{=} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Since this reduced matrix lies in \mathfrak{N} we don't need any compression and get the following corresponding tableau

$$S(M_1) = \begin{array}{|c|c|}\hline 1\\\hline 2\\\hline 2&3\\\hline \end{array}.$$

For M_2 we write:

$$Y_2(3)^{-2} = X_3(2)^2 X_4(1)^2,$$

 $Y_3(2)^{-1} = X_4(2),$
 $Y_1(1)^3 = X_1(1)^3,$
 $Y_1(3) = X_1(3).$

Hence

$$M_2 \leftrightarrow \begin{pmatrix} 3 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 2 & 0 \\ 2 & 1 & 0 \end{pmatrix}.$$

One step of compression yields

$$M_1^{(1)} = \begin{pmatrix} 3 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 2 & 0 \\ 3 & 0 & 0 \end{pmatrix} \leftrightarrow S(M_2) = \underbrace{\begin{array}{c} 1 \\ \hline 1 & 1 & 1 & 4 & 4 & 4 \end{array}}_{\boxed{1 & 1 & 1 & 4 & 4 & 4}}.$$

Due to the reversed bumping rule in [8] we get

On the other hand we observe for the monomial bumping

$$M_1 * M_2 = \begin{pmatrix} & & 0 & & & \\ & & 0 & & & \\ & M_2 & \vdots & & M_1 & \\ & & 0 & & & \\ & & 0 & & & \end{pmatrix} = \begin{pmatrix} 3 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 2 & 0 & 0 & 1 & 0 & 0 \\ 2 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and by compression we get

$$(M_1 * M_2)^{(3)} = \begin{pmatrix} 3 & 0 & 1 & 1 \\ 0 & 0 & 2 & 0 \\ 0 & 2 & 0 & 0 \\ 3 & 0 & 0 & 0 \end{pmatrix} \stackrel{(A1)}{=} \begin{pmatrix} 3 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 0 \end{pmatrix}.$$

Hence we get the expected equation:

Chapter 3

Type C

3.1 Compression of Nakajima monomials in type C

In this section we will define the compression of Nakajima monomials for a Lie algebra \mathfrak{g} of type C_n . We shortly recall the basic setting of \mathfrak{g} . Let P be the weight lattice of \mathfrak{g} and β_1, \ldots, β_n the orthogonal basis of P. Let further $I = \{1, \ldots, n\}$ be the index set for the simple roots given by $\alpha_i = \beta_i - \beta_{i+1}$ for $i = 1, \ldots, n-1$ and $\alpha_n = 2\beta_n$. Moreover we get the fundamental weights by $\Lambda_i = \beta_1 + \ldots + \beta_i$ and therefore $\beta_i = \Lambda_i - \Lambda_{i-1}$. Then we compute for all $i \neq n$ and $j \in I$:

$$\langle h_j, \alpha_i \rangle = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } j = i - 1 \text{ or } j = i + 1, \\ 0 & \text{else} \end{cases}$$

and

$$\langle h_j, \alpha_n \rangle = \begin{cases} 2 & \text{if } j = n, \\ -2 & \text{if } j = n - 1, \\ 0 & \text{else.} \end{cases}$$

As in the A_n -case we set

$$c_{ij} = \begin{cases} 0 & \text{if } i > j, \\ 1 & \text{else.} \end{cases}$$

With this notation we obtain for $j \in \mathbb{Z}$

$$A_i(j) = \begin{cases} Y_i(j)Y_{i+1}(j)^{-1}Y_i(j+1)Y_{i-1}(j+1)^{-1} & \text{if } i \neq n, \\ Y_n(j)Y_n(j+1)Y_{n-1}(j+1)^{-2} & \text{if } i = n. \end{cases}$$

Let $\mathbf{B} = \{1, \dots, n, \bar{1}, \dots \bar{n}\}$ then we define a total order on \mathbf{B} by

$$1 \prec 2 \prec \ldots \prec n \prec \overline{n} \prec \ldots \prec \overline{2} \prec \overline{1}$$
.

For $i \in I$ and $j \in \mathbb{Z}$ we consider the variables defined in [3]:

$$X_i(j) := Y_{i-1}(j+1)^{-1}Y_i(j),$$

 $X_{\overline{i}}(j) := Y_{i-1}(j+n-i+1)Y_i(j+n-i+1)^{-1}.$

With these variables we have for $i \neq n$:

$$A_i(j) = X_{i+1}(j)^{-1}X_i(j),$$

 $A_i(j) = X_{\overline{i+1}}(j-n+i)X_{\overline{i}}(j-n+i)^{-1}$

and

$$A_n(j) = X_n(j)X_{\overline{n}}(j)^{-1}.$$

Furthermore it is easy to see that for $i=1,\ldots,n$ and some p-q=n-i the following equation holds

$$X_i(p)X_{\overline{i}}(q) = X_{i+1}(p)X_{\overline{i+1}}(q).$$

This equation will be important later when we define the equivalence relation on matrices because it involves more options to write an arbitrary monomial as a product of $X_i(j)$'s and $X_{\bar{i}}(j)$'s.

As in Chapter 2 we recall the characterization of $M_1(\lambda)$ with a dominant integral weight λ for Lie algebras of type C_n given in [3]:

Proposition 3.1 [3] Let $\lambda = a_1\Lambda_1 + \ldots + a_n\Lambda_n$. Then the connected component $M_1(\lambda)$ containing the maximal vector

$$M_1 = Y_1(1)^{a_1} \cdots Y_n(1)^{a_n}$$

is characterized as the set of monomials

$$M = X_{t_{1,1}}(1) \cdots X_{t_{1,k_1}}(1) \cdots X_{t_{n,1}}(n) \cdots X_{t_{n,k_n}}(n)$$

satisfying the following conditions:

(i)
$$k_i = a_i + \ldots + a_n$$
 for all $j = 1, \ldots, n$,

(ii)
$$t_{j,1} \succeq t_{j,2} \succeq \ldots \succeq t_{j,k_j}$$
 for all $j = 1, \ldots, n$,

(iii) for each
$$j = 2, ..., n$$
 and $l = 1, ..., k_j, t_{j-1,l} > t_{j,l}$.

For $s \in \mathbb{Z}$ we also consider the *shifted* highest weight monomials of weight λ

$$M_s = Y_1(s)^{a_1} Y_2(s)^{a_2} \dots Y_n(s)^{a_n}.$$

As an immediate consequence of Proposition 3.1 we obtain their connected component $\mathcal{M}_s(\lambda)$ by the set of monomials of the form

$$M = X_{t_{1,1}}(s) \cdots X_{t_{1,k_1}}(s) \cdots X_{t_{n,1}}(s+n-1) \cdots X_{t_{n,k_n}}(s+n-1)$$

satisfying condition (i), (ii) and (iii).

We will see later on that these conditions translate into the notation of matrices exactly the same way as in the A_n -case. In order to use similar constructions as in Chapter 2 we show that \mathfrak{M} is generated by the elements $X_i(j)$ and $X_{\bar{i}}(j)$.

That means we define **M** to be the free abelian monoid generated by the set $\{X_i(j), X_{\overline{i}}(j), i = 1, \dots, n, j \in \mathbb{Z}\}$. We further define an ideal $\mathbf{J} \subset \mathbf{M}$ by

$$\mathbf{J} := \langle \prod_{k=1}^{i} X_k(j+i-k) X_{\overline{k}}(j-n+k-1), \\ \prod_{k=1}^{i} X_{\overline{k}}(j-i+k) X_k(j+n+1-k), i = 1, \dots, n, j \in \mathbb{Z} \rangle_{\mathbf{M}}.$$

Hence the quotient M/J becomes a group with

$$X_i(j)^{-1} = \prod_{k=1}^{i-1} X_k(j+i-k) \prod_{k=1}^{i} X_{\overline{k}}(j-n+k-1)$$

and

$$X_{\overline{i}}(j)^{-1} = \prod_{k=1}^{i-1} X_{\overline{k}}(j-i+k) \prod_{k=1}^{i} X_k(j+n+1-k).$$

This gives rise to an analog of Proposition 2.2.

Proposition 3.2 We have

$$\mathfrak{M} \cong \mathbf{M}/\mathbf{J}$$
.

Proof: We consider the map that identifies $X_i(j)$ with $Y_i(j)Y_{i-1}(j+1)^{-1}$ and $X_{\overline{i}}(j)$ with $Y_{i-1}(j+n-i+1)Y_i(j+n-i+1)^{-1}$. Let $M=\prod_{i\in I, j\in \mathbb{Z}}Y_i(j)^{y_i(j)}$

be a monomial in \mathfrak{M} . In order to show surjectivity we consider again each $Y_i(j)^{y_i(j)}$ separately and distinguish two cases:

1. case: $y_i(j) > 0$, then we write

$$Y_{i}(j) = \prod_{k=0}^{i-1} Y_{k}(j+i-k)^{-1} \prod_{k=1}^{i} Y_{k}(j+i-k)$$
$$= \prod_{k=1}^{i} X_{k}(j+i-k).$$

Therefore we get

$$Y_i(j)^{y_i(j)} = \prod_{k=1}^{i} X_k(j+i-k)^{y_i(j)}.$$

2. case: $y_i(j) < 0$, then we set

$$Y_i(j)^{-1} = \prod_{k=1}^{i} X_{\overline{k}}(j-n+k-1)$$

and hence

$$Y_i(j)^{y_i(j)} = \prod_{k=1}^i X_{\overline{k}}(j-n+k-1)^{-y_i(j)}.$$

With these equations we compute

$$X_i(j)^{-1} = Y_i(j)^{-1} Y_{i-1}(j+1) = \prod_{k=1}^{i-1} X_k(j+i-k) \prod_{k=1}^{i} X_{\overline{k}}(j-n+k-1)$$

and

$$X_{\overline{i}}(j)^{-1} = Y_{i-1}(j+n-i+1)^{-1}Y_i(j+n-i+1)$$

=
$$\prod_{k=1}^{i-1} X_{\overline{k}}(j-i+k) \prod_{k=1}^{i} X_k(j+n+1-k),$$

which implies injectivity.

Remark 3.1 The equation

$$X_i(p)X_{\overline{i}}(q) = X_{i+1}(p)X_{\overline{i+1}}(q)$$

for p - q = n - i, also holds in \mathbf{M}/\mathbf{J} .

Due to Proposition 3.2 we can write an arbitrary $M \in \mathfrak{M}$ as a product of $X_i(j)$'s and $X_{\overline{i}}(j)$'s. More precisely, there exist $m_{ij} \in \mathbb{Z}_{\geq 0}$ such that

$$M = \prod_{i \in \mathbf{B}, j \in \mathbb{Z}} X_i(j)^{m_{ij}}.$$

In other words, we can associate a matrix $m_{i,j}$ to each $M \in \mathfrak{M}$, where $i \in$ $\{1,\ldots,n,1,\ldots\bar{n}\}\ \mathrm{and}\ j\in\mathbb{Z}.$ That means we obtain a matrix in $\mathrm{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0}).$ As in the A_n -case these are matrices with just finitely many non zero columns and we number the rows by $1, \ldots, n, \bar{n}, \ldots \bar{1}$ instead of $1, \ldots, 2n$.

In order to get a bijection between the monomials and those matrices we need to fix the matrix notation. Consider $M \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{>0})$ with $M = m_{i,j}$. Then the definition of J and Remark 3.1 allow us to apply the following rules without changing the underlying monomial:

(C1) For a pair p, q with p - q = n - b with

$$m_{b,p} \neq 0$$
 and $m_{\bar{b},q} \neq 0$

we decrease $m_{b,p}$ and $m_{\overline{b},q}$ by $\min\{m_{b,p}, m_{\overline{b},q}\}$ and increase $m_{b+1,p}$ and $m_{\overline{b+1},q}$ by $\min\{m_{b,p}, m_{\overline{b},q}\}$.

(C2) For a pair p, q with p - q = n - b + 1 with

$$m_{b,p} \neq 0$$
 and $m_{\bar{b},q} \neq 0w$

we decrease $m_{b,p}$ and $m_{\overline{b},q}$ by $\min\{m_{b,p},m_{\overline{b},q}\}$ and increase $m_{b-1,p}$ and $m_{\overline{b-1},q}$ by $\min\{m_{b,p},m_{\overline{b},q}\}$.

Moreover we have an analog of rule (A1):

(C3) For every $i \in \{1, ..., n\}$ and $k \in \mathbb{Z}$ with

$$m_{i-s,k+s} \neq 0$$
 for all $s = 0, 1, ..., i-1$

and

$$m_{\overline{i-s},k-n+i-1-s} \neq 0$$
 for all $s=0,1,\ldots,i-1$

we decrease all these entries by

$$\min\{m_{i-s,k+s}, m_{\overline{i-s},k-n+i-1-s}; 0, 1, \dots, i-1\}.$$

We call such a collection a generalized diagonal at $m_{i,k}$ and this procedure a cancellation at $m_{i,k}$.

We can also insert some generalized diagonals to get longer ones:

(C4) For every $i \in \{1, ..., n\}$ and $k \in \mathbb{Z}$ with

$$m_{i,k} \neq 0$$

and

$$m_{\overline{i-s},k-n+i-1-s} \neq 0$$
 for all $s = 0,1,\ldots,i-1$

we increase the entries $m_{i-s,k+s}$ for all $s=1,\ldots,i-1$ by

$$\min\{m_{i,k}, m_{\overline{i-s},k-n+i-1-s}; 0, 1, \dots, i-1\}$$

and apply (C3) to get a longer cancellation at $m_{i,k}$.

For every $i \in \{1, ..., n\}$ and $k \in \mathbb{Z}$ with

$$m_{\bar{i},k-n+i-1} \neq 0$$

and

$$m_{i-s,k+s} \neq 0$$
 for all $s = 0, 1, \dots, i-1$

we increase the entries $m_{\overline{i-s},k-n+i-1-s}$ for all $s=1,\ldots,i-1$ by

$$\min\{m_{i-s,k+s}, m_{\bar{i},k-n+i-1}; 0, 1, \dots, i-1\}$$

and apply (C3) to get a longer cancellation at $m_{i,k}$.

We use the rules (C1)-(C4) to get reduced versions of the matrices associated to monomials in \mathfrak{M} . We explain what we mean by reduced in this case: Let $m_{i,j} \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ be a matrix corresponding to a monomial $M \in \mathfrak{M}$:

$$M = \prod_{i \in \mathbf{B}, j \in \mathbb{Z}} X_i(j)^{m_{ij}}.$$

We search for $[m_{i,j}]$ such that

(i)
$$\sum_{i \in \mathbf{B}, j \in \mathbb{Z}} [m_{ij}] = \min \{ \sum_{i \in \mathbf{B}, j \in \mathbb{Z}} n_{ij} ; \prod_{i \in \mathbf{B}, j \in \mathbb{Z}} X_i(j)^{m_{ij}} = \prod_{i \in \mathbf{B}, j \in \mathbb{Z}} X_i(j)^{n_{ij}} \},$$

(ii) there are no pairs p, q with p - q = n - b + 1 such that

$$[m_{b,p}] \neq 0$$
 and $[m_{\overline{b},q}] \neq 0$.

We use the rules (C1)-(C4) stepwise to obtain $[m_{ij}]$ from m_{ij} as follows:

Let j be minimal such that $m_{i,k} = 0$ for all $i \in \mathbf{B}$ and k > j. Then we start at $m_{n,j}$ and apply (C1)-(C2) to all the other entries if this yields an application of (C3)-(C4) to $m_{n,j}$. This means we try to get some cancellation at this entry. After that we go left to the next entry in this row and do the same.

Once we have done this with the whole row we go to the upper one and apply the same procedure until we reach $m_{1,1}$.

At the end we apply (C2) to guarantee the desired condition (ii).

Example 3.1 For \mathfrak{g} of type C_3 we consider the monomial

$$M = Y_1(0)Y_1(2)Y_1(1)^{-1}Y_1(5)^{-1}Y_1(3)^{-1}Y_1(4)^{-2}Y_2(0)Y_2(3)Y_2(5)^{-2}Y_3(0)Y_3(4).$$

Due to Proposition 3.1 we write

$$\begin{array}{rcl} Y_1(0) & = & X_1(0), \\ Y_1(2) & = & X_1(2), \\ Y_1(1)^{-1} & = & X_{\overline{1}}(-2), \\ Y_1(5)^{-1} & = & X_{\overline{1}}(2), \\ Y_1(3)^{-1} & = & X_{\overline{1}}(0), \\ Y_1(4)^{-2} & = & X_{\overline{1}}(1)^2, \\ Y_2(0) & = & X_2(0)X_1(1), \\ Y_2(3) & = & X_2(3)X_1(4), \\ Y_2(5)^{-2} & = & X_{\overline{2}}(3)^2X_{\overline{1}}(2)^2, \\ Y_3(0) & = & X_3(0)X_2(1)X_1(2), \\ Y_3(4) & = & X_3(4)X_2(5)X_1(6) \end{array}$$

and assign the matrix

We observe that we get no (C3) or (C4) for $m_{3,j}$. That means we go to i = 2 and get a generalized diagonal at $m_{2,5}$. After applying cancellation we obtain the new martix by

The next entry we have to consider is $m_{2,4}$ and after insertion at $m_{1,5}$ and application of (C1) we also get a generalized diagonal at $m_{2,4}$ and hence

where we omit the zero-columns. In the next step we apply (C1) to $m_{1,0}$ and insert a generalized diagonal at $m_{1,0}$ to get a cancellation at $m_{2,0}$, which yields the reduced version matrix

$$\begin{pmatrix} 1 & 0 & 2 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 \end{pmatrix} = [m_{i,j}].$$

With this notation we define an equivalence relation on $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$:

$$m_{i,j} \sim n_{i,j} \text{ iff } [m_{i,j}] = [n_{i,j}].$$

We consider the quotient

$$\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{>0})/\sim.$$

It is obvious that two matrices which lie in the same equivalence class correspond to the same monomial and vice versa. Hence by sending a monomial onto its reduced matrix we get

Proposition 3.3 There exists a bijection between \mathfrak{M} and $Mat_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$.

In order to get a morphism of crystals we endow $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ with a crystal structure by defining it on the reduced representatives.

Let $M = m_{i,j} \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ be a reduced matrix. Then we set

$$wt(M) = \sum_{i=1}^{n} \left(\sum_{j \in \mathbb{Z}} (m_{ij} - m_{\bar{i},j}) \right) \beta_i.$$

For $i \neq n$ we put

$$\varphi_{i}(M) = \max \left\{ \sum_{j \leq k} m_{ij} + m_{\overline{i+1},j-n+i} - \sum_{j < k} m_{i+1,j} + m_{\overline{i},j-n+i} ; k \in \mathbb{Z} \right\},
\epsilon_{i}(M) = -\min \left\{ \sum_{j > k} m_{ij} + m_{\overline{i+1},j-n+i} - \sum_{j \geq k} m_{i+1,j} + m_{\overline{i},j-n+i} ; k \in \mathbb{Z} \right\}$$

and

$$\varphi_n(M) = \max \left\{ \sum_{j \le k} m_{n,j} - \sum_{j < k} m_{\overline{n},j} ; k \in \mathbb{Z} \right\},$$

$$\epsilon_n(M) = -\min \left\{ \sum_{j > k} m_{n,j} - \sum_{j \ge k} m_{\overline{n},j} ; k \in \mathbb{Z} \right\}.$$

If $\varphi_i(M) = 0$ we set $\tilde{f}_i(M) = 0$ for all $i \in I$.

Let now $\varphi_i(M) \neq 0$ then we define the Kashiwara operator \tilde{f}_i for $i \neq n$: Let k be minimal such that

$$\varphi_i(M) = \sum_{j \le k} m_{ij} + m_{\overline{i+1},j-n+i} - \sum_{j < k} m_{i+1,j} + m_{\overline{i},j-n+i}.$$

Then we distinguish the following two cases:

<u>1. case</u>: $m_{\overline{i+1},k-n+i} = 0$.

Then we set $\tilde{f}_i(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{i+1,k}$ (resp. $m_{i,k}$) by one. Formally spoken we obtain $\tilde{f}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(i,k), (i+1,k)\}, \\ m_{i,k} - 1 & \text{if } (s,j) = (i,k), \\ m_{i+1,k} + 1 & \text{if } (s,j) = (i+1,k). \end{cases}$$

 $\underline{2. \text{ case}}: m_{\overline{i+1},k-n+i} \neq 0.$

Then we define $\tilde{f}_i(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{\bar{i},k-n+i}$ (resp. $m_{\bar{i}+1,k-n+i}$) by one. That means we obtain $\tilde{f}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(\bar{i},k-n+i), (\bar{i+1},k-n+i)\}, \\ m_{\bar{i}+1,k-n+i} - 1 & \text{if } (s,j) = (\bar{i+1},k-n+i), \\ m_{\bar{i},k-n+i} + 1 & \text{if } (s,j) = (\bar{i},k-n+i). \end{cases}$$

Now we give the definition of $\tilde{f}_n(M)$ for $\varphi_n(M) \neq 0$: Let k be minimal such that

$$\varphi_n(M) = \sum_{j \le k} m_{n,j} - \sum_{j \le k} m_{\overline{n},j}.$$

Then we set $\tilde{f}_n(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{\overline{n},k}$ (resp. $m_{n,k}$) by one. More precisely we obtain $\tilde{f}_n(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(n,k), (\overline{n},k)\}, \\ m_{n,k} - 1 & \text{if } (s,j) = (n,k), \\ m_{\overline{n},k} + 1 & \text{if } (s,j) = (\overline{n},k). \end{cases}$$

If $\epsilon_i(M) = 0$ we set $\tilde{e}_i(M) = 0$.

For $\epsilon_i(M) \neq 0$ let p be maximal such that

$$\epsilon_i(M) = -(\sum_{j>p} m_{ij} + m_{\overline{i+1},j-n+i} - \sum_{j\geq p} m_{i+1,j} + m_{\overline{i},j-n+i}).$$

Then we distinguish the following two cases to define $\tilde{e}_i(M)$ for $i \neq n$:

1. case: $m_{\overline{i+1},p-n+i} \neq 0$.

Then we set $\tilde{e}_i(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{i,p}$ (resp. $m_{i+1,p}$) by one. Formally spoken we observe $\tilde{e}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(i,p), (i+1,p)\}, \\ m_{i,p} + 1 & \text{if } (s,j) = (i,p), \\ m_{i+1,p} - 1 & \text{if } (s,j) = (i+1,p). \end{cases}$$

<u>2. case</u>: $m_{\overline{i+1},n-n+i} = 0$.

Then we defin $\tilde{e}_i(M)$ as the matrix we get from M by increasing (resp. decreasing) $m_{\overline{i+1},p-n+i}$ (resp. $m_{\overline{i},p-n+i}$) by one. That means we obtain $\tilde{e}_i(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(\overline{i},p-n+i), (\overline{i+1},p-n+i)\}, \\ m_{\overline{i+1},p-n+i} + 1 & \text{if } (s,j) = (\overline{i+1},p-n+i), \\ m_{\overline{i},p-n+i} - 1 & \text{if } (s,j) = (\overline{i},p-n+i). \end{cases}$$

Let p be maximal such that

$$\epsilon_n(M) = -(\sum_{j>p} m_{n,j} - \sum_{j>p} m_{\overline{n},j}).$$

Then we set $\tilde{e}_n(M)$ as the matrix we get from M by decreasing (resp. increasing) $m_{\overline{n},p}$ (resp. $m_{n,p}$) by one. Formally spoken we obtain $\tilde{e}_n(M) = \hat{m}_{s,j}$ from $M = m_{s,j}$ by

$$\hat{m}_{s,j} = \begin{cases} m_{s,j} & \text{if } (s,j) \notin \{(n,p), (\overline{n},p)\}, \\ m_{n,p} + 1 & \text{if } (s,j) = (n,p), \\ m_{\overline{n},p} - 1 & \text{if } (s,j) = (\overline{n},p). \end{cases}$$

Easy computations show that $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ along with the maps $wt, \varphi_i, \epsilon_i, \tilde{f}_i$ and \tilde{e}_i becomes a semi-normal crystal. As in section 2 we prove that this crystal structure coincides with the structure on \mathfrak{M} under the above bijection.

Proposition 3.4 The bijection

$$\Psi: \qquad \mathfrak{M} \qquad \to Mat_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$$

$$M = \prod_{i \in \mathbf{B}, j \in \mathbb{Z}} X_i(j)^{m_{ij}} \qquad \mapsto [m_{ij}]$$

is a crystal isomorphism.

Proof: It is easy to verify that wt, ϵ_i and φ_i are invariant under Ψ especially because they are invariant under the application of (C1)-(C4). It remains to show that Ψ commutes with the crystal operators \tilde{f}_i and \tilde{e}_i . Let $M \in \mathfrak{M}$ be a

monomial with associated matrix $[m_{l,j}]$. Due to the crystal structure defined on the matrices it follows almost directly that for all $i \in I$

$$\tilde{f}_i(M) = \prod_{l \in \mathbf{B}, j \in \mathbb{Z}} X_l(j)^{\tilde{f}_i([m_{lj}])}$$

and the same for \tilde{e}_i .

Therefore it suffices to verify that

$$\tilde{f}_i([m_{lj}]) = [\tilde{f}_i([m_{lj}])]$$

and

$$\tilde{e}_i([m_{lj}]) = [\tilde{e}_i([m_{lj}])].$$

Since this can be proved analogously we just give the proof for \tilde{f}_i .

For simplicity we denote $[m_{l,j}]$ by $m_{l,j}$ and for $i \neq n$ let k be minimal such that

$$\varphi_i(M) = \sum_{j \le k} m_{ij} + m_{\overline{i+1},j-n+i} - \sum_{j < k} m_{i+1,j} + m_{\overline{i},j-n+i}.$$

First we look at the case $m_{\overline{i+1},k-n+i}=0$. That means we get $\tilde{f}_i(m_{l,j})$ by increasing $m_{i+1,k}$ and decreasing $m_{i,k}$ each by one. Let us assume that the increase of $m_{i+1,k}$ induces a longer or new cancellation. But this yields

$$m_{i+1,k} = 0$$
 and $m_{\bar{i},k-n+i} = 0$

and together with the choice of k this implies

$$m_{i,k+1} = 0$$
 and $m_{\overline{i+1},k-n+i+1} = 0$.

Since $m_{i,k+1} = 0$ and $m_{\tilde{i},k-n+i} = 0$, the increase doesn't provide any generalized diagonal at $m_{i+s,k+1-s}$ for positive s, without insertion. But those insertions would have been done before we operate with \tilde{f}_i because $m_{l,j}$ is in reduced form.

It is still possible that we get a new generalized diagonal by (C4) at $m_{i+1,k}$ itself. But we get no diagonal $m_{i+1-s,k+s} \neq 0$ for $s = 0, \ldots, i$ since $m_{i,k+1} = 0$. Furthermore we can't increase this entry by applying (C1) because otherwise we could have applied (C4) at $m_{i,k+1}$ before.

The second possibility to apply (C4) at $m_{i+1,k}$ needs $m_{\overline{i+1-s},k-n+i-s} \neq 0$ for all $s = 0, \ldots, i$. In particular we get $m_{\overline{i-s},k-n-1+i-s} \neq 0$ for all $s = 0, \ldots, i-1$ and $m_{i,k} \neq 0$ which implies an application of (C4) at $m_{i,k}$ before operating. This is again a contradiction to the fact that $m_{i,j}$ is reduced.

Moreover since $m_{\overline{i+1},k-n+i} = 0$ we can not apply (C1) to the increased $m_{i+1,k}$. Overall we have seen that operating with \tilde{f}_i preserves the reduced version in this case.

Similar arguments hold for the case $m_{\overline{i+1},k-n+i} \neq 0$.

Now we translate the characterizing conditions of the monomials that give a realization of the crystal bases, given in Proposition 3.1, into the language of matrices. We will recognize that those are the same conditions as in Chapter 2. From this observation one can deduce that similar constructions yield our desired morphism.

Definition 3.1 Define $\mathfrak{N} \subset Mat_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ as the set of matrices whose reduced versions have only zero-entries out of a $2n \times n$ -submatrix $M = (m_{ij})_{\substack{i=1,\ldots,n,\overline{n},\ldots,\overline{1}\\j=0,\ldots,n-1}}$ satisfying the following properties:

(i)
$$m_{ij} \in \mathbb{Z}_{>0}$$
 for $i = 1, \ldots, n, \overline{n}, \ldots, \overline{1}$ and $j = 0, \ldots, n-1$,

(ii)
$$\sum_{k\geq i} m_{k,j} \leq \sum_{k>i} m_{k,j-1}$$
 for $i=1,\ldots,n,\overline{n},\ldots,\overline{1}$ and $j=1,\ldots,n-1,$ where we set $\sum_{k>i} m_{k,j-1} = 0$ for $i=\overline{1}$.

Due to the crystal structure and the equivalence relation above we observe the following remark which helps us to guarantee that operating interchanges with lower decomposition later. Moreover it implies that \mathfrak{N} and hence also \mathfrak{N} are stable under application of \tilde{e}_i and \tilde{f}_i .

Remark 3.2 Let $m_{i,j}$ be a reduced version of a matrix in $Mat_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})$ and $i\in I$.

- (i) If \tilde{f}_i acts on $m_{i,k}$ then $m_{i,k} > m_{i+1,k-1}$,
- (i) if \tilde{e}_i acts on $m_{i+1,p}$ then $m_{i+1,p} > m_{i,p+1,p}$

where we set
$$i + 1 = \overline{i-1}$$
 if $i \in \{\overline{n}, \dots, \overline{2}\}$ and $n + 1 = \overline{n}$.

As mentioned above we also use the lower decomposition rule for the later constructions. Therefore we need an C-analog of Lemma 2.1.

Lemma 3.1 Let $[M] = (m_{ij})_{i=1,\ldots,n,\overline{n},\ldots,\overline{1}} \in Mat_{2n\times l}(\mathbb{Z}_{\geq 0})$ be a reduced matrix without zero columns which satisfies condition (ii) of Definition 3.1. Then we have

$$l < n$$
.

Proof: Assume l > n and consider a special collection of elements in $\mathbf{B} = \{1, \dots, n, \bar{n}, \dots \bar{1}\}$:

For
$$k = 1, ..., n + 1$$
 let $i_k \in \mathbf{B}$ be maximal such that $m_{i_k, k} \neq 0$.

This collection exists because there are no zero columns and l > n. Furthermore condition (ii) of Definition 3.1 implies

$$i_{n+1} \prec i_n \prec \ldots \prec i_2 \prec i_1$$
.

That means there exists at least one pair $p, q \in \{1, \dots, n+1\}$ with p > q such that

$$i_p \in \{1, \dots, n\}$$
 and $i_q = \overline{i_p}$.

Let p be minimal with this property. The minimality of p yields

$$p-q \leq n-i_p+1$$
.

We assume $p-q < n-i_p+1$, namely $p-q=n-i_p+1-j$ for some $j \in \mathbb{N}$. Let us consider the number of elements between i_p and i_q :

$$|\{i_p, i_{p-1}, \dots, i_{q+1}, i_q\}| = n - i_p - j + 2.$$

Since

$$(n - i_p - j + 2) + (i_p - 1) = n - j + 1 < n + 1$$

there is another pair \hat{p} , \hat{q} with $\hat{p} > p > \hat{q}$ such that $i_{\hat{p}} \in \{1, \dots, n\}$ and $i_{\hat{q}} = \overline{i_{\hat{p}}}$. If we consider \hat{p} minimal with this property one gets

$$\hat{p} - \hat{q} \le n - i_{\hat{p}} + 1.$$

If we assume $\hat{p} - \hat{q} < n - i_{\hat{p}} + 1$ we can use the same arguments as above. This way we can inductively conclude that there has to be such a pair with $p - q = n - i_p + 1$ which is a contradiction to $m_{i,j}$ being reduced and therefore $l \leq n$.

Let us define the C-analog of the map Φ given in Section 2.1.

$$\Phi : \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0}) / \sim \to \mathfrak{N} / \sim$$

$$m_{ij} \mapsto n_{ij},$$

where we compute n_{ij} as follows:

Let M be a reduced version matrix in $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$. Then we consider the lower decomposition of M:

$$M = M_1 + M_2,$$

with $M_1 = m_{i,j}^{(1)}$, $M_2 = m_{i,j}^{(2)}$ and $M_1 \in \mathfrak{N}$. We use exactly the same decomposition as in Section 2.1 with 2n rows instead of n+1. Then we move every entry of M_2 one column to the left and denote this matrix by $M_2^{(1)}$ and set

$$M^{(1)} := M_1 + M_2^{(1)}.$$

Then we decompose $[M^{(1)}]$ and proceed the same way until the iteration becomes stationary and we reach $M^{(k)}=m_{i,j}^{(k)}\in\mathfrak{N}$. Then set

$$n_{i,j} = N := M^{(k)}.$$

Before we show that this map has the desired properties we state another lemma which will be useful for the proof of the main theorem.

Lemma 3.2 Let $M=M_1+M_2$ be the lower decomposition of a matrix in reduced form with $M_1=m_{i,j}^{(1)}$ and $M_2=m_{i,j}^{(2)}$. Then there exists no pair p,q with p-q=n-i such that

$$m_{i,p}^{(1)} \neq 0 \text{ and } m_{\bar{i},q}^{(2)} \neq 0.$$

Proof: Due to the lower decomposition rule we obtain $M_1 \in \mathfrak{N}$ and since M is reduced it's obvious that M_1 is reduced. Now one can use the same arguments as used in the proof of Lemma 3.1.

We define the compression map κ again as the following composition

$$\kappa := \Psi^{-1} \circ \Phi \circ \Psi$$

and show

37

Theorem 3.1 Let \mathfrak{g} be of type C. Then, the map

$$\kappa : \mathfrak{M} \to \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$$
$$M \mapsto (\Psi^{-1} \circ \Phi \circ \Psi)(M)$$

defined as above is a morphism of crystals.

Proof: We limit ourselves to prove that sending a reduced matrix M onto $M^{(1)}$ thus defined preserves the crystal structure. This implies inductively that Φ and hence $\Psi^{-1} \circ \Phi \circ \Psi$ are crystal morphisms. So consider $M = m_{l,j}$ the reduced form of an arbitrary matrix in $\operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ and $m_{l,j} = m_{l,j}^{(1)} + m_{l,j}^{(2)}$ its lower decomposition. For an $i \in I$ we have to show:

(i)
$$wt(M) = wt(M^{(1)}),$$

(ii)
$$\varphi_i(M) = \varphi_i(M^{(1)}),$$

(iii)
$$\epsilon_i(M) = \epsilon_i(M^{(1)})$$

and that computing $M^{(1)}$ commutes with the Kashiwara operators namely

(iv)
$$(\tilde{f}_i(M))^{(1)} = \tilde{f}_i(M^{(1)}),$$

(v)
$$(\tilde{e}_i(M))^{(1)} = \tilde{e}_i(M^{(1)}).$$

We confine ourselves to prove (ii) and (iv) because the rest follows analogously. Rather we just show (ii) and get (iv) from the A_n -case. Set $N = M_1 + M_2^{(1)}$ with $N = n_{l,j}$ and let k be minimal such that

$$\varphi_i(M) = \sum_{j \le k} m_{ij} + m_{\bar{i}+1,j-n+i} - \sum_{j < k} m_{i+1,j} + m_{\bar{i},j-n+i}.$$

In the first step we show $\varphi_i(M) \leq \varphi_i(N)$. For simplicity we introduce some notation:

For $i \in I, l \in \mathbb{Z}$ and a matrix $M = m_{i,j}$ we set

$$p_{i,l}(M) := \sum_{j \le l} m_{i,j} + m_{\overline{i+1},j-n+i} - \sum_{j < l} m_{i+1,j} + m_{\overline{i},j-n+i}.$$

First of all we look at the case $m_{\overline{i+1},k-n+i}=0$. That yields $m_{i,k}\neq 0$ and furthermore Remark 3.2 says $m_{i,k}>m_{i+1,k-1}$. The fact that M is reduced also implies $m_{\overline{i},k-n+i-1}=0$. We show that either

$$p_{i,k}(N) = p_{i,k}(M) = \varphi_i(M)$$

or

$$p_{i,k-1}(N) = p_{i,k}(M) = \varphi_i(M).$$

Let us assume that $p_{i,k}(N) < p_{i,k}(M)$. The first case that could yield this is

$$\sum_{j \le k} n_{i,j} - \sum_{j < k} n_{i+1,j} < \sum_{j \le k} m_{i,j} - \sum_{j < k} m_{i+1,j}.$$

But the computation in the proof of Theorem 2.1 shows that in this case we obtain

$$\sum_{j \le k} m_{i,j} - \sum_{j < k} m_{i+1,j} = \sum_{j \le k-1} n_{i,j} - \sum_{j < k-1} m_{i+1,j}$$

and since $m_{\overline{i+1},k-n+i}=0, m_{\overline{i},k-n+i-1}=0$ we get

$$\varphi_i(M) = p_{i,k-1}(N).$$

The other possibility to get $p_{i,k}(N) < p_{i,k}(M)$ is

$$\sum_{j \le k} n_{i,j} - \sum_{j < k} n_{i+1,j} = \sum_{j \le k} m_{i,j} - \sum_{j < k} m_{i+1,j}$$

and

$$\sum_{j \leq k} n_{\overline{i+1},j-n+i} - \sum_{j \leq k} n_{\overline{i},j-n+i} < \sum_{j \leq k} m_{\overline{i+1},j-n+i} - \sum_{j \leq k} m_{\overline{i},j-n+i}.$$

The first equation implies $0 \neq m_{i,k}^{(1)} > m_{i+1,k-1}^{(1)}$ and the inequation yields

 $0 \neq m_{\bar{i},k-n+i}^{(2)} > m_{\bar{i}+1,k-n+i+1}^{(2)}$. But the existence of $0 \neq m_{i,k}^{(1)}$ and $0 \neq m_{\bar{i},k-n+i}^{(2)}$ provides a contradiction to Lemma 3.2.

Now we consider the case $m_{\overline{i+1},k-n+i} \neq 0$. Since M is reduced we get $m_{i+1,k} = 0$ and in particular $m_{i+1,k}^{(2)} = 0$. Therefore the only chance to have $p_{i,k}(N) < p_{i,k}(M)$ is $m_{i,k}^{(1)} \neq 0$ and $0 \neq m_{\overline{i},k-n+i}^{(2)} > m_{\overline{i+1},k-n+i+1}^{(2)}$, which is again a contradiction to Lemma 3.2.

Overall we get

$$\varphi_i(M) < \varphi_i(N)$$
.

It remains to show:

$$\varphi_i(M) \ge \varphi_i(N)$$
.

Suppose $\varphi_i(M) < \varphi_i(N)$, that means there is a $t \in \mathbb{Z}$ with

$$p_{i,t}(N) > \varphi_i(M)$$
.

We distinguish the same cases as above. At first we assume

$$\sum_{j \le t} n_{i,j} - \sum_{j \le t} n_{i+1,j} > \sum_{j \le t} m_{i,j} - \sum_{j \le t} m_{i+1,j}.$$

That is only possible if $m_{i,t+1}^{(2)} > m_{i+1,t}^{(2)}$. In particular we obtain $m_{i,t+1} \neq 0$ and hence $m_{\bar{i},p-n+i} = 0$. From the A_n -case we know that the following inequation holds in this case

$$\sum_{j \le t} n_{i,j} - \sum_{j < t} n_{i+1,j} \le \sum_{j \le t+1} m_{i,j} - \sum_{j < t+1} m_{i+1,j}.$$

Combining this with $m_{\bar{i},p-n+i}=0$ and $p_{i,t}(N)>\varphi_i(M)$ we see

$$p_{i,t+1}(M) > \varphi_i(M),$$

which is a contradiction to the choice of k. Now suppose that

$$\sum_{j \le t} n_{\overline{i+1},j-n+i} - \sum_{j < t} n_{\overline{i},j-n+i} > \sum_{j \le t} m_{\overline{i+1},j-n+i} - \sum_{j < t} m_{\overline{i},j-n+i}.$$

That means $0 \neq m_{\overline{i+1},t-n+i+1}^{(2)} > m_{\overline{i},t-n+i}^{(2)}$ and the A_n -case implies again

$$\sum_{j < t} n_{\bar{i}, j - n + i} - \sum_{j < t} n_{\bar{i}, j - n + i} \le \sum_{j < t + 1} m_{\bar{i}, j - n + i} - \sum_{j < t + 1} m_{\bar{i}, j - n + i}.$$

If $m_{i,p+1} = 0$ we obtain the same contradiction as in the above case. So we assume $m_{i,p+1} \neq 0$. But in order to get $p_{i,t}(N) > \varphi_i(M)$ we need $m_{i,p+1}^{(1)} \neq 0$ because otherwise

$$p_{i,t}(N) \le p_{i,t+1}(M) \le \varphi_i(M).$$

But $m_{i,p+1}^{(1)} \neq 0$ and $m_{\overline{i+1},t-n+i+1}^{(2)} \neq 0$ provide a contradiction to Lemma 3.2. Hence

$$\varphi_i(M) \ge \varphi_i(N)$$
.

Moreover these arguments also show that either k or k-1 is minimal such that either

$$\varphi_i(N) = p_{i,k}(N)$$

or

$$\varphi_i(N) = p_{i,k-1}(N).$$

Finally Remark 3.2 and the A_n -case imply (iv) which finishes our proof.

Example 3.2 For n = 3 we reconsider Example 3.1 namely the monomial

$$M = Y_1(0)Y_1(2)Y_1(1)^{-1}Y_1(5)^{-1}Y_1(3)^{-1}Y_1(4)^{-2}Y_2(0)Y_2(3)Y_2(5)^{-2}Y_3(0)Y_3(4).$$

We have already seen that we can write M as

$$M = X_1(0)X_1(2)X_{\overline{1}}(-2)X_{\overline{1}}(2)X_{\overline{1}}(0)X_{\overline{1}}(1)^2X_2(0)X_1(1)X_2(3)X_1(4)X_{\overline{2}}(3)^2 X_{\overline{1}}(2)^2X_3(0)X_2(1)X_1(2)X_3(4)X_2(5)X_1(6)$$

with reduced version matrix

$$[m_{i,j}] = \begin{pmatrix} 1 & 0 & 2 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 \end{pmatrix}.$$

40

We observe its lower decomposition by

and therefore

$$[m_{i,j}]^{(1)} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 2 & 2 & 0 & 0 \end{pmatrix} = [[m_{i,j}]^{(1)}].$$

One further step yields the desired matrix in \mathfrak{N} :

$$\Phi(\Psi(M)) = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 4 & 0 & 0 \end{pmatrix}$$

and by application of Ψ^{-1} we get the monomial

$$N = X_1(0)X_1(1)X_1(2)X_2(0)X_2(1)^2X_2(2)X_{\overline{2}}(0)X_{\overline{3}}(0)^4$$

= $Y_1(0)Y_2(0)Y_2(1)^2Y_1(3)^{-1}Y_3(1)^{-4}Y_2(1)^4 \in \mathcal{M}_0(3\Lambda_1 + 2\Lambda_2 + 2\Lambda_3).$

Let B(M) be the connected component of $M \in \mathfrak{M}$.

Corollary 3.1 For $M \in \mathfrak{M}$ and $\Phi(\Psi(M)) =: n_{i,j} \in \mathfrak{N}$ we consider $s \in \mathbb{Z}$ maximal such that $n_{i,j} = 0$ for all j < s and $i \in \{1, \ldots, n, \overline{n}, \ldots \overline{1}\} = \mathbf{B}$. Furthermore for $k = 1, \ldots, n$ we define the values:

$$a_k := \sum_{i \in B} n_{i,k+s-1} - \sum_{i \in B} n_{i,k+s} \ge 0.$$

Then we have

$$\kappa(M) \in \mathcal{M}_s(\sum_{k=1}^n a_k \Lambda_k)$$

and by restricting our morphism to the connected component we get that

$$\kappa_{|B(M)}$$
 : $B(M) \rightarrow \mathcal{M}_s(\sum_{k=1}^n a_k \Lambda_k)$

is a crystal isomorphism.

3.2 Monomials and C-Tableaux

Kim and Shin [7] also gave a realization of the crystal bases in the sense of reversed Young tableaux for Lie algebras of type C. In this case they obtained $S(\lambda)$ as the set of all semistandard reversed Young tableaux of shape λ with entries $1, \ldots, n, \overline{n}, \ldots, \overline{1}$ satisfying some conditions (for details see [7]). Moreover Kang, Kim and Shin [3] constructed a morphism between $\mathcal{M}_1(\lambda)$ and those tableaux for \mathfrak{g} of type C_n . This is similar to the one in Section 2.2 and can also be generalized to a crystal morphism between arbitrary monomials and tableaux in $S(\lambda)$ via compression.

Let M be in $\mathcal{M}_1(\lambda)$ for an integral dominant weight λ and $m_{i,j}$ its associated reduced matrix. For $i \in \mathbf{B}$ we define again S(M) to be the reversed tableaux with $m_{i,j}$ many i's in the j-th row. We will say that there is an i(p) if there exists an entry i in the p-th row of the tableaux. In order to get a tableaux that satisfies the condition of $S(\lambda)$ we have to apply the following rules which are due to [3]:

(al-1) For each $a=1,\ldots,n-1$, if there is a pair $(a(p),\overline{a}(q))$ such that p-q< n-a and either a(p) and $\overline{a}(q)$ lie in the same column or $\overline{a}(q)$ lies in the left hand side of a(p).

Then replace $(a(p), \overline{a}(q))$ with $(a + 1(p), \overline{a + 1}(q))$.

If there are several such pairs then we begin with the rightmost a(p) and leftmost $\overline{a}(q)$.

(al-2) For each b = n, ..., 2, if there is a pair $(b(p), \overline{b}(q))$ such that p - q = n - b + 1 and $\overline{b}(q)$ lies in the right hand side of b(p).

Then replace $(b(p), \overline{b}(q))$ with $(b-1(p), \overline{b-1}(q))$.

If there are several such pairs then we begin with the leftmost b(p) and rightmost $\bar{b}(q)$.

Let us denote by [S(M)] the reversed tableaux we obtain from S(M) by applying those rules. Then we can state the following

Proposition 3.5 [3] The map

$$\begin{array}{cccc} \Omega & : & M_1(\lambda) & \to & S(\lambda) \\ & & M & \mapsto & [S(M)] \end{array}$$

is a crystal isomorphism.

As in the A_n -case we continue this morphism to $\mathcal{M}_s(\lambda)$. Let $M \in \mathcal{M}_s(\lambda)$ be a monomial and m_{ij} the associated reduced matrix in \mathfrak{N} . We set S(M) to be the semistandard reversed tableaux with m_{ij} -many i entries in the j-s+1-st row and get the morphism

$$\Omega : \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda) \to \bigcup_{\lambda \in P} S(\lambda)$$

$$M \mapsto [S(M)].$$

If we combine this result with Theorem 3.1 we get a morphism between Nakajima monomials and tableaux:

Corollary 3.2 The map

$$\Omega\circ\kappa \ : \ \mathfrak{M} \ \to \ \bigcup_{\lambda\in P}S(\lambda)$$

is a crystal morphism.

Let us consider an example:

Example 3.3 For \mathfrak{g} of type C_3 consider the monomial

$$M = Y_2(2)^2 Y_2(1)^{-1} Y_3(0) Y_1(0) Y_3(3)^{-1}.$$

Via compression we get $\Phi(\Psi(M))$ by

$$\begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

This yields the following tableau:

$$S(\kappa(M)) = \underbrace{ \begin{array}{c|c} \boxed{1} \\ 2 \ \hline 3 \\ \hline \end{array} }_{ \begin{array}{c|c} \boxed{1} \ 2 \ \hline \end{array} } \underbrace{ \begin{array}{c|c} \boxed{2} \\ \hline \hline \end{array} }_{ \begin{array}{c|c} \boxed{1} \ 2 \ \hline \end{array} } \underbrace{ \begin{array}{c|c} \boxed{3} \\ \hline \hline \end{array} }_{ \begin{array}{c|c} \boxed{3} \ \hline \hline \end{array} } = [S(\kappa(M))] \in S(\lambda).$$

3.3 Insertion scheme for monomials in type C

In this section we define a C-analogue of the bumping rule for Nakajima monomials given in Section 2.3. Let $M_1, M_2 \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ be two reduced version matrices of monomials in \mathfrak{M} . As in the A_n -case we need to associate a matrix $M_1 * M_2 \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0})$ to $M_1 \otimes M_2$. In order to assure that $M_1 * M_2$ is in reduced form we need more zero columns between M_1 and M_2 in this case. Namely we insert n zero-columns and define

$$M_1 * M_2 = \begin{pmatrix} & 0 & \dots & 0 \\ & 0 & & 0 \\ & M_2 & \vdots & & \vdots & M_1 \\ & 0 & & 0 \\ & 0 & \dots & 0 \end{pmatrix} \in \operatorname{Mat}_{2n \times \mathbb{Z}}(\mathbb{Z}_{\geq 0}).$$

With the tensor product rule and the same arguments as in Proposition 2.6 we observe:

Proposition 3.6 The map

$$\begin{array}{cccc} \operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \otimes \operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim & \to & \operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \\ M_1\otimes M_2 & \mapsto & M_1*M_2 \end{array}$$

is a crystal morphism.

Furthermore for $M_1, M_2 \in \mathfrak{M}$ we can define the bumping $M_2 \to M_1$ via compression analogously to the A_n -case:

$$M_1 \to M_2 := \Psi^{-1}(\Phi(\Psi(M_1) * \Psi(M_2))),$$

where Ψ is the crystal isomorphism between \mathfrak{M} and $\operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim$ and $\Phi: \operatorname{Mat}_{2n\times\mathbb{Z}}(\mathbb{Z}_{\geq 0})/\sim \mathfrak{N}/\sim$ the matrix compression. Theorem 3.1 and Proposition 3.6 imply

Theorem 3.2 Let \mathfrak{g} be of type C. Then, the map

$$\mathfrak{M} \otimes \mathfrak{M} \quad \to \quad \bigcup_{\lambda \in P, s \in \mathbb{Z}} \mathcal{M}_s(\lambda)$$
 $M_1 \otimes M_2 \quad \mapsto \quad M_1 \to M_2$

is a crystal morphism.

Kim and Shin [7],[8] also defined a bumping rule for reversed tableaux in type C. Therefore it is natural to compare $[S(M_1 \to M_2)]$ and $[S(M_1)] \to [S(M_2)]$ as in Section 2.3, where $[S(M_1 \to M_2)]$, $[S(M_1)]$ and $[S(M_2)]$ are the corresponding tableaux in $\bigcup_{\lambda} S(\lambda)$ due to Corollary 3.2. Theorem 3.2 together with Corollary 3.2 imply again

$$[S(M_1 \to M_2)] = [S(M_1)] \to [S(M_2)].$$

Example 3.4 For \mathfrak{g} of type C_3 we compute the bumping of the two monomials

$$M_1 = Y_3(0)Y_3(2)Y_2(1)Y_2(3)^{-2}Y_1(3)Y_1(2)^{-1}$$

and

$$M_2 = Y_2(2)^2 Y_2(1)^{-1} Y_3(0) Y_1(0) Y_3(3)^{-1}.$$

At first we consider the monomial bumping $M_1 \to M_2$ and its associated tableau $[S(M_1 \to M_2)]$. Afterwards we compute the tableaux bumping of $[S(M_1)]$ and $[S(M_2)]$, namely $[S(M_1)] \to [S(M_2)]$. Then we will obtain the expected equation

$$[S(M_1 \to M_2)] = [S(M_1)] \to [S(M_2)].$$

We compute the reduced version matrices of M_1 and M_2 by

$$\Psi(M_2) = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \Psi(M_1) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 0 \end{pmatrix}.$$

Therefore we have to apply the compression map Φ to the matrix

We apply the compression stepwise and get

$$(M_1 * M_2)^{(1)} = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(1)}],$$

$$(M_1 * M_2)^{(2)} = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(2)}],$$

$$(M_1 * M_2)^{(3)} = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 2 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 2 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(3)}],$$

$$(M_1 * M_2)^{(4)} = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 2 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 2 & 0 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(4)}],$$

$$(M_1 * M_2)^{(5)} = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 2 & 0 & 0 \end{pmatrix} \Rightarrow [(M_1 * M_2)^{(5)}] = \begin{pmatrix} 1 & 0 & 2 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 \end{pmatrix},$$

$$(M_1 * M_2)^{(6)} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 2 & 2 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(6)}]$$

and finally

$$(M_1 * M_2)^{(7)} = egin{pmatrix} 1 & 1 & 1 \ 1 & 2 & 0 \ 1 & 0 & 1 \ 0 & 0 & 0 \ 0 & 1 & 0 \ 4 & 0 & 0 \end{pmatrix} = [(M_1 * M_2)^{(7)}] \in \mathfrak{N}.$$

Hence we observe the corresponding reversed tableau

$$S(M_1 \to M_2) = \frac{\begin{array}{c|c} 1 & 3 \\ \hline 1 & 2 & 2 & \overline{2} \\ \hline 1 & 2 & 3 & \overline{1} & \overline{1} & \overline{1} \end{array}}{\begin{array}{c|c} 1 & 2 & 3 \\ \hline \end{array}} \in S(3\Lambda_1 + 2\Lambda_2 + 2\Lambda_3)$$

On the other hand we assign the tableaux to M_1 and M_2 separately:

By applying the reversed bumping rule for tableaux we get

$$[S(M_1)] \to [S(M_2)] = \begin{array}{c|c} \hline 3 \\ \hline 2 & 2 & \overline{2} \\ \hline 3 & \overline{1} & \overline{1} \end{array} \to \begin{array}{c|c} \hline 2 \\ \hline 3 & \overline{3} \\ \hline 1 & 2 & \overline{3} & \overline{3} \\ \hline \hline 2 & \overline{3} & \overline{3} & \overline{2} \end{array}$$
$$= \begin{array}{c|c} \hline 2 & 3 \\ \hline 1 & 2 & 3 & \overline{2} \\ \hline \hline 1 & 2 & \overline{3} & \overline{1} & \overline{1} & \overline{1} \\ \hline \end{array}$$
$$= [S(M_1 \to M_2)].$$

References

- [1] Kang, S.-J., Kim, S.-J., Lee, H., Shin, D.-U.: Young wall realization of crystal bases for classical Lie algebras. Trans. Amer. Math. Soc. **356**, 2349-2378 (2004)
- [2] Kang, S.-J., Kim, S.-J. Shin, D.-U.: Monomial realization of crystal bases for special linear Lie algebras. J. Algebra 274, 629-642 (2004)
- [3] Kang, S.-J., Kim, J.-A., Shin, D.-U.: Crystal bases for quantum classical algebras and Nakajima's monomials. Publ. Res. Inst. Math. Sci. 40, 758-791 (2004)
- [4] Kashiwara, M.: On crystal bases of the q-analogue of universal enveloping algebras. Duke Math. J. **63**, 465-516 (1991)
- [5] Kashiwara, M.: Realizations of crystals. Contemp. Math. 325, 133-139 (2003)
- [6] Kashiwara, M., Nakashima, T.: Crystal graphs for representations of the q-analogue of classical Lie algebras. J. Algebra 165, 295-345 (1994)
- [7] Kim, J.-A., Shin, D.-U. Correspondence between Young walls and Young tableaux realizations of crystal bases for the classical Lie algebras. J. Algebra 282, 728-757 (2004)
- [8] Kim, J.-A., Shin, D.-U. Insertion schemes for the classical Lie algebras. Communications in Algebra **32**, 3139-3167 (2004)
- [9] Littelmann, P. Paths and root operators in representation theory. Annals of Math. 142, 449-525 (1995)
- [10] Littelmann, P. Characters of representations and paths in $\mathfrak{h}_{\mathbb{R}}^*$. Proceedings of Symposia in Pure Mathematics **61**, 29-49 (1997)
- [11] Nakajima H., t-analogs of q-characters of quantum affine algebras of type A_n and D_n . Contemp. Math. **325**, 141-160 (2003)

Erklärung

Ich versichere, dass ich die von mir vorgelegte Dissertation selbständig angefertigt, die benutzten Quellen und Hilfsmittel vollständig angegeben und die Stellen der Arbeit - einschliesslich Tabellen, Karten und Abbildungen -, die anderen Werken im Wortlaut oder dem Sinn nach entnommen sind, in jedem Einzelfall als Entlehnung kenntlich gemacht habe; dass diese Dissertation noch keiner anderen Fakultät oder Universität zur Prüfung vorgelegen hat; dass sie - abgesehen von unten angegebenen Teilpublikationen - noch nicht veröffentlicht worden ist sowie, dass ich eine solche Veröffentlichung vor Abschluss des Promotionsverfahrens nicht vornehmen werde. Die Bestimmungen dieser Promotionsordnung sind mir bekannt. Die von mir vorgelegte Dissertation ist von Prof. Dr. Littelmann betreut worden.

keine Teilpublikationen -	
Köln, den 20. November 2010	
	(Matthias Meng)

Lebenslauf

Angaben zur Person

Nachname, Vorname | Meng, Matthias

Staatsangehörigkeit Deutsch

Geburtsdatum 5. November 1983

Geschlecht männlich

Geburtsort Sangerhausen

Familienstand ledig, keine Kinder

Studium

vorauss. 02/2011 Abschluss der Promotion

Titel der Dissertation:

"Compression of Nakajima monomials in type A and C"

seit 05/2009 Promotionsstipendium im Graduiertenkolleg "Globale Strukturen in Geometrie

und Analysis" an der Universität zu Köln

Forschungsbereich: Kombinatorische Darstellungstheorie

Betreuer: Prof. Dr. Peter Littelmann

03/2009 Diplom in Mathematik mit Nebenfach Informatik

Abschlussnote: sehr gut

Titel der Diplomarbeit: Nakajima-Monome und Wegemodell (Ausgezeichnet mit

einem Hauptpreis der DMV-Studierendenkonferenz 2009)

Köln

03/2007 Vordiplom in Mathematik mit Nebenfach Informatik

Note: sehr gut

10/2004 - 03/2007 | Grundstudium der Mathematik mit Nebenfach Informatik an der Universität

zu Köln

Studienbegleitende Tätigkeiten

seit 08/2005 Pädagogische Mitarbeit bei der Übermittagsbetreuung der Königin-Luise-Schule

Köln

04/2009 Tätigkeit als wissenschaftliche Hilfskraft am Lehrstuhl Prof. Dr. Peter Littel-

nann

10/2005 - 03/2009 Tätigkeit als Übungsgruppenleiter in den Fächern Lineare Algebra, Stochastik

und Zahlentheorie an der Universität zu Köln

Auslandsaufenthalte

08/2010 2-wöchiger Aufenthalt in Nagoya, Japan

Besuch der Konferenz:

"Representation Theory of Algebraic Groups and Quantum Groups"

09/2009 1-wöchiger Aufenthalt in Ascona, Schweiz

Besuch der Sommerschule:

"Algebraic Groups and Invariant Theory"

07/2009 3-wöchiger Aufenthalt in Ottawa, Kanada

Besuch der Sommerschule und Konferenz:

"Geometric Representation Theory and Extended Affine Lie Algebras"

Fort- und Weiterbildungen

11/2009 "Präsentation - Sicher auftreten und gekonnt reden"

Sabine Hahn, Theater Akademie Köln

Zivildienst

08/2003 - 05/2004 | Im Kinder- und Jugendzentrum Neheim

Schulbildung

08/1994 - 06/2003 Abitur am St.-Ursula Gymnasium Neheim

Note: 1,8

08/1991 - 06/1994 Besuch der Karl-Wagenfeld Grundschule Neheim

08/1990 - 06/1991 Besuch der Bernard-Koenen Oberschule Sangerhausen

Fremdsprachenkenntnisse

Englisch sicher in Wort und Schrift

Französisch Grundkenntnisse

EDV-Kenntnisse

 $\begin{array}{ccc} \text{MS-Office} & \text{sehr gute Kenntnisse} \\ \text{I\!AT}_{E\!X} & \text{sehr gute Kenntnisse} \end{array}$

Java Grundkenntnisse