

Supplementary material

Plant host and microbe specific gene expression in the mutualistic root endophytic fungus *Serendipita vermifera*

Mahdi LK^{*1}, Eichfeld R^{*1}, Miyauchi S^{*2}, Concetta De Quattro¹, Langen G¹, Vivian Ng³, Daniel Peterson³, Igor Grigoriev³, Yu Zhang, Sravanthi³, Tejomurhula³ and Zuccaro, A¹.

¹ Institute of Plant Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), Cologne Biocenter, University of Cologne, Zùlpicher StraÙe 47a, 50674 Cologne, Germany.

² Okinawa Institute of Science and Technology Graduate University (OIST), 1919-1 Tancha, Onna, Kunigami District, Okinawa 904-0495 Japan.

³Joint Genome Institute – Lawrence Berkeley Laboratory, 1 Cyclotron Road Berkeley, CA 94720 USA

* These authors contributed equally to this work.

Author for correspondence: Alga Zuccaro, azuccaro@uni-koeln.de

Supplementary methods

Cloning: To clone *SvCHIT-CBM5* and *SiCHIT-CBM5* without signal peptide into the empty vector (Fig. S5) for expression in *E. coli*, both sequences were amplified from cDNA using the program and primers listed in Table S1+2. PCR products run on 1 % agarose gels. Bands were extracted and purified using the NucleoSpin® Gel and PCR Clean-up kit (Macherey-Nagel). The concentration of the PCR product was adjusted to a three-fold molar excess of plasmid. The plasmid was linearized with restriction enzyme KpnI and cloning was performed via Gibson Assembly (Table S3). The reactions incubated at 50 °C for 15 min. The resultant Plasmids as pictured in Fig. S6+7.

Table S1: PCR program for the amplification of *SvCHIT-CBM5* and *SiCHIT-CBM5* including the individual steps, temperature, and duration

| Step | Temperature (°C) | Time (mm:ss) |
|----------------------|------------------|--------------|
| Initial denaturation | 95 | 00:30 |
| Denaturation | 95 | 00:20 |
| Primer annealing | 62 | 00:20 |
| Extension | 72 | 00:30 per Kb |
| Final extension | 72 | 02:00 |
| Hold | 4 | ~ |

Table S2: Primer list for the amplification of *SvCHIT-CBM5* and *SiCHIT-CBM5* listing all primer-names and sequences

| Name | Sequence |
|-------------------------------|--|
| <i>SvCHIT-CBM5</i> (Forward) | ATCACCATCACCATCACGGATCCGCATGCG AGCTCGGTACCACTCCCGTCATGCAACGAC |
| <i>SvCHIT -CBM5</i> (Reverse) | CTCAGCTAATTAAGCTTGGCTGCAGGTCGA CCCGGGGTACCTCAGCAGGCTCGCACCT |
| <i>SiCHIT -CBM5</i> (Forward) | ATCACCATCACCATCACGGATCCGCATGCG AGCTCGGTACCAACGCCCGCCATGATGC |
| <i>SiCHIT-CBM5</i> (Reverse) | CTCAGCTAATTAAGCTTGGCTGCAGGTCGA CCCGGGGTACCTCAGCACGACTTGAGTTGA TTCCAC |

Table S3: Gibson Assembly mix showing the components and amounts, that vary depending on the Plasmid concentration and fragment length and was calculated via the formula below.

| Component | Amount |
|----------------------------|--------------------|
| Plasmid | X μ L |
| PCR fragment | Y μ L |
| 2 x Gibson Master Mix | 10 μ L |
| Deionized H ₂ O | 10 – (X+Y) μ L |

$$\frac{\text{Plasmid concentration in ng}/\mu\text{L} * 1000}{\text{length of fragment in bp} * 650 \text{ kDa}} = \text{pmol}/\mu\text{L}$$

Transformation of *E. coli*: 10 μ L plasmid were added to competent *E. coli* cells (Mach1) and incubated for 20 min on ice. Plasmid uptake was stimulated by a 45 sec incubation at 42 °C and subsequent incubation on ice for 2 min. The cells regenerated in 500 μ L LS-LB medium at 37 °C for at least 30 min under continuous shaking, where then centrifuged for 30 sec at 11,000 rpm. The cells were resuspended in 20 μ L LS-LB medium, plated on carbenicillin containing LS-LB agar plates and incubated at 37 °C overnight. Colony PCR components and program are listed in Table S4.

Table S4: Colony PCR Components

| Component | Amount for 25 μ L reaction | |
|---------------------------|--------------------------------|--|
| 10 x Thermo pol buffer | 2.5 μ L | |
| 10 mM dNTPs | 0.5 μ L | |
| Forward primer | 0.5 μ L | |
| Reverse primer | 0.5 μ L | |
| <i>Taq</i> DNA polymerase | 0.125 μ L | |
| H ₂ O | To 25 μ L | |

| Step | Temperature (°C) | Time (mm:ss) |
|----------------------|------------------|--------------|
| Initial denaturation | 98 | 00:30 |
| Denaturation | 98 | 00:30 |
| Primer annealing | 45-68 | 00:30 |
| Extension | 68 | 01:00 per kB |
| Final extension | 68 | 05:00 |
| Hold | 4 | ~ |

PCR products were run on 1 % agarose gels to confirm successful transformation of cloned plasmids.

Mini- and Maxi plasmid DNA preparation: Mini- and Maxi plasmid DNA preparations were performed from either 4 mL or 300 mL bacteria cultures grown in LS-LB medium containing the appropriate antibiotics after incubation at 37°C and 180 rpm using either the NucleoSpin^R Plasmid kit (Macherey & Nagel) or the NucleoBond^R Xtra Maxi kit (Macherey & Nagel). Plasmid concentration was measured via nanodrop.

SDS-PAGE and Western Blot: *E. coli* crude extracts were measured by Bradford assay and equal protein concentrations were adjusted. The supernatant was boiled in SDS loading dye for 5 min at 95 °C. 20 µL of sample were loaded on 10 x SDS-Gels and run at 190 V for 50 min. Proteins were blotted on PVDF membranes with transfer buffer for 1 h at 17 V. The membrane was blocked in TBS-T-BSA overnight at 4 °C and next the HisProbe-HRP conjugate was added (1:20000 dilution) was added in TBS-T-BSA and incubated 1 h at RT. Next, the membrane was washed 3 times in TBS-T and 3 times in TBS (each washing step 5 min). Developing solution was added to detect chemiluminescence

Supplementary figures




Figure S1: Distribution and density of the normalised \log_2 transformed read counts. At: *Arabidopsis thaliana*; Bd: *Brachypodium distachyon*; Hv: *Hordeum vulgare*; Sv: *Serendipita vermifera*; Bs: *Bipolaris sorokiniana*; Bact: bacteria; d: days a) Before removal of the inconsistent samples AtSv_3d. b) After removal of AtSv_3d. A set of data excluding AtSv_3d was used for co-expression and DEG analyses.




Figure S2: Correlations of RNA-seq Samples. Left part: Adjacent matrix of the Spearman correlation coefficients. Right part: Hierarchical clustering of biological replicates based on distances of transcriptomic similarities. At: *Arabidopsis thaliana*; Bd: *Brachypodium distachyon*; Hv: *Hordeum vulgare*; Sv: *Serendipita vermifera*; Bs: *Bipolaris sorokiniana*; Bact: bacteria; d: days.




Figure S3: Differentially expressed genes (DEGs) of *Sv* in response to different interaction partners and timepoints. UpSetplot of significantly differentially expressed genes (DEGs) (FDR-adjusted $p < 0.05$) compared to control (fungus alone). Dotted lines indicate a combination of conditions. *At*: *Arabidopsis thaliana*; *Bd*: *Brachypodium distachyon*; *Hv*: *Hordeum vulgare*; *Sv*: *Serendipita vermifera*; *Bs*: *Bipolaris sorokiniana*; Bact: bacteria; dpi: days post inoculation

